



**“.. I think this is maybe our Achilles Heel....” Exploring GPs’ responses to young people presenting with emotional distress in general practice. A qualitative study.**

Journal:	<i>BMJ Open</i>
Manuscript ID:	bmjopen-2013-002927
Article Type:	Research
Date Submitted by the Author:	21-Mar-2013
Complete List of Authors:	Roberts, Jane; University of Sunderland, Pharmacy, Health & Well-being Crosland, Ann; University of Sunderland, Pharmacy, Health & Well-being Fulton, John; University of Sunderland, Pharmacy, Health & Well-being
<b>Primary Subject Heading</b>:	General practice / Family practice
Secondary Subject Heading:	Mental health, Medical education and training, Qualitative research
Keywords:	PRIMARY CARE, MENTAL HEALTH, Child & adolescent psychiatry < PSYCHIATRY

SCHOLARONE™  
Manuscripts

1  
2  
3  
4  
5 **“.. I think this is maybe our Achilles Heel....” Exploring GPs’ responses to young people presenting**  
6 **with emotional distress in general practice. A qualitative study.**  
7  
8  
9

10  
11  
12 Jane H Roberts , Ann Crosland, John Fulton

13  
14 Department of Pharmacy, Health & Well-being, Faculty of Applied Sciences, The Science Complex,  
15 Wharnccliffe St, Sunderland, SR1 3SD.  
16

17  
18 Jane H Roberts, Clinical Senior Lecturer, University of Sunderland,  
19

20  
21 Ann Crosland, Professor of Nursing , University of Sunderland,  
22

23  
24 John Fulton, Principal Lecturer, University of Sunderland,  
25

26  
27 Correspondence to Jane H Roberts  
28

29  
30 jane.roberts@sunderland.ac.uk  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3  
4  
5 **“.. I think this is maybe our Achilles Heel....” Exploring GPs’ responses to young**  
6 **people presenting with emotional distress in general practice. A qualitative study.**  
7  
8  
9

## 10 **Article summary**

### 11 **Article Focus:**

- 12 1. an exploratory study
- 13 2. to examine GPs’ views and experiences of consulting with young people
- 14 3. to understand GPs perspectives

### 15 **Key Messages**

- 16 1. GPs collectively describe anxiety and uncertainty about their clinical
- 17 2. practice when consulting with young people in distress, independently of age and
- 18 3. gender
- 19 2. Anxiety relates to professional performance; interacting with young people
- 20 3. and the complex nature of presentations of emotional distress in primary care
- 21 3. Unless anxiety and uncertainty are addressed GPs will continue to miss
- 22 4. opportunities to address early emotional difficulties and young people’s mental
- 23 5. health needs in primary care will continue to be poorly met

### 24 **Strengths and Limitations**

- 25 1. Qualitative research in under -examined areas offers new insights and
- 26 2. explores why behaviours might arise
- 27 3. The data contributes to theory building and offers theoretical
- 28 4. generalizability
- 29 5. Theoretical sampling led to only white British born GPs participating so
- 30 6. other cultural perspectives were not included

1  
2  
3 “.. I think this is maybe our Achilles Heel....” Exploring GPs’ responses to young  
4 people presenting with emotional distress in general practice. A qualitative study.  
5  
6  
7  
8

## 9 Main text

### 10 Introduction

11  
12  
13  
14 Emotional distress in young people is common. It may indicate an associated mental  
15 health problem, with at least 10% of 10-15 year olds affected <sup>1</sup> and 17 % of 16-19  
16 year olds <sup>2</sup> (based on household surveys). Proxy markers of distress, such as  
17 reported incidences of self-harm derived from community based studies, show 10%  
18 of adolescents report having self-harmed. <sup>3</sup>  
19

20  
21  
22  
23 Data from populations of young people who consult their GP reveal higher rates of  
24 psychological distress, of the order of 20-30%. <sup>4 5</sup> GPs identify serious mental illness  
25 but often fail to detect less severe manifestations <sup>6</sup> and appear reluctant to discuss  
26 emotional issues<sup>7</sup>; unless offered cues by the young person in the consultation <sup>8</sup> or if  
27 other factors are present such as a previous history of a suicide attempt or a pattern  
28 of frequent consulting <sup>9</sup>. Young people’s presentations in primary care are often  
29 complex and present with behavioural, psychosocial, academic and familial  
30 problems which can be problematic to untangle. They may suggest underlying co-  
31 morbid mental health problems. It has been reported that often the ‘most important  
32 features in terms of assessment may be concealed or hidden’. <sup>10</sup>  
33  
34

35  
36  
37  
38 A key concern is the difficulty of distinguishing between ‘moodiness’ or a persisting  
39 emotional disorder and GPs have expressed a worry at ‘over-medicalising young  
40 people’s lives’. <sup>11</sup> Illiffe & colleagues found that GPs were uncomfortable about  
41 making a diagnosis of depression in young people (the most common, but often  
42 coexisting, mental health problem in adolescence ).  
43  
44  
45  
46  
47  
48

49  
50 This sits in contrast to GPs’ increasing involvement of common mental health  
51 problems in older patients <sup>12 13</sup> and also to a broadening of the frames of reference  
52 by which emotional distress in adults is regarded. Although a biomedical perspective  
53 dominates, supported by an array of NICE clinical guidelines, Dowrick <sup>14</sup> and Reeve  
54 <sup>15</sup> have offered alternative frameworks and refer to the insights derived from the  
55  
56  
57  
58  
59  
60

1  
2  
3 wisdom traditions. Historically, GPs have been found to be dismissive of their role in  
4 addressing social issues in adult mental ill-health<sup>16</sup> although this position is shifting  
5 with greater awareness of the lay perspective, which typically favours the causes of  
6 mental ill-health ( notably depression) as being social in origin<sup>17</sup>.  
7  
8  
9

10 Despite the challenge of responding to emotional distress in adolescence and the  
11 patchy, often inadequate provision of secondary care services<sup>18 19</sup> a series of policy  
12 directives have emphasised the role of GPs and other front-line services, in the  
13 promotion of psychological well-being and the early indication of difficulties.<sup>20 21 22</sup>  
14 Practitioners are expected to have 'sufficient knowledge, training and support 'in this  
15 area including competence in 'active listening' and conversational technique'<sup>23</sup>.  
16  
17  
18  
19

20 There is a growing body of evidence examining young people's experiences of  
21 talking to GPs about emotional problems. They reveal a mixed picture including a  
22 reluctance to disclose<sup>24</sup>, a fear of being judged or offered medication<sup>25</sup>. Much less  
23 is known about GP perspectives. This paper presents a qualitative, exploratory study  
24 which examines GPs' views and experiences of consulting with young people  
25 presenting with emotional distress.  
26  
27  
28  
29  
30  
31

## 32 **Method**

### 33 *Study Design*

34  
35 The study took place in the North East of England in 18 general practices based in  
36 urban, rural and semi-rural communities serving predominantly socio-economically  
37 disadvantaged patients. The qualitative study comprised of in depth individual  
38 interviews with GPs recruited using theoretical sampling. As early theoretical ideas  
39 emerged successive GPs were recruited on the basis of their capacity to contribute  
40 to the development or abandonment of initial theoretical constructs.  
41  
42  
43  
44  
45  
46

47 Data were collected between January 2010 to May 2011  
48  
49

### 50 *Participants*

51  
52 GPs with less than four years clinical experience were excluded. The initial recruits  
53 were selected on the basis of their relevant experience and their ability to generate  
54 early data which would scope the terrain of the area under enquiry.  
55  
56  
57  
58  
59  
60

1  
2  
3 GPs were approached by telephone and email contact and sent information sheets.  
4 A follow-up contact established their verbal consent to meet at a location of their  
5 choice.  
6  
7

### 8 9 *Data collection and analysis*

10  
11 The audio-taped semi-structured interviews were transcribed verbatim with consent.  
12 An initial topic guide was used with the first tranche of participants based on the  
13 extant literature and developed through discussion. The topic guide was then revised  
14 on the basis of ideas arising from the early interviews, and the iterative analysis  
15 which began as soon as the first interview was undertaken. The interview guides  
16 explored doctors' experiences of consulting with young people in general and those  
17 presenting with psychological or mental health problems, GPs' understanding of  
18 depression and anxiety in adolescence, of how emotional distress presents in the  
19 surgery and the role of the GP in promoting emotional well-being in young people  
20 (See appendix 1). The guide was refined to include questions about how structural  
21 changes impacted on, and consultation style shaped, practice.  
22  
23  
24  
25  
26  
27  
28

29  
30 The interviews lasted between 50 to 75 minutes. Field notes and theoretical memos  
31 were kept throughout the period of data collection and analysis.  
32  
33

34 The transcripts were coded and analysed using the grounded theory method  
35 described by Strauss and Glaser<sup>26</sup> and revised by Charmaz.<sup>27</sup> The constant  
36 comparative method of analysis is core to the process and informs the theoretical  
37 sampling of recruits. Situational maps, both 'messy' and 'ordered', were constructed  
38 during this phase of analysis.<sup>28</sup>  
39  
40  
41  
42

43 The data presented here was produced after the first level of analysis was completed  
44 during which the open codes were developed by JR and subject to further  
45 examination by AC (primary care academic) and JF(sociologist)  
46  
47  
48

### 49 **Results**

50  
51 Nineteen GPs participated, 10 women. (Table1). The early iterative analysis of the  
52 data found a dominant narrative of anxiety and uncertainty about practice under-  
53 pinning the majority of the research interviews. This pervasive and disabling  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 emotional response to encounters with emotionally distressed young people  
4 appeared to coalesce around three domains.  
5  
6

7 These can be viewed as anxiety and uncertainty experienced by GPs in response to:  
8

9  
10 1) professional performance; in the consultation, at an external level, across  
11 disciplinary boundaries;  
12

13  
14 2) interacting with young people; and  
15

16  
17 3) the complexity of presentations of adolescent emotional distress  
18

19 **Anxiety related to professional performance: *In the consultation***  
20

21 A prevailing finding was the sense of professional impotence which was associated  
22 with seeing or suspecting emotional distress in this age group. It was acknowledged  
23 that feeling unsure of practice led to a sense of disempowerment through not  
24 knowing what to do; in contrast to working with older patients where the options  
25 appear more clearly defined. The data collected suggested that not being able to  
26 formulate the initial presentation by a young person into a definable 'disorder'  
27 created a sense of operating in uncharted territory.  
28  
29

30  
31 This was amplified by the lack of exposure to adolescent mental health in  
32 undergraduate medical education which was unanimously shared by all participants.  
33  
34 Where the topic had been included in the curriculum, it was often restricted to severe  
35 mental disorder such as adolescents hospitalized with anorexia nervosa.  
36  
37

38  
39 **Anxiety related to professional performance: *at an external level***  
40

41 The lack of benchmarks meant assessing one's performance in relation to peers was  
42 problematic since no 'gold standard' existed. The only NICE guideline which was  
43 referenced (concerning the management of depression in under 18 year olds) was  
44 regarded as having hampered GPs from becoming involved in the management of  
45 adolescent depression and supporting a view that there was little to be offered in  
46 primary care.  
47  
48

49  
50 Varying arrangements within practices governing access to appointments and the  
51 ease, or not, of maintaining continuity of care were seen to contribute to professional  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 anxiety by impeding attentive 'watchful waiting' and some GPs described attempts to  
4 circumvent inflexible appointment systems in order to be more available to patients.  
5  
6

7 A lack of professional supervision was identified by a small number of more  
8 experienced GPs involved with Postgraduate Training and provision of mental  
9 health services at a regional level, and contrasted to systems for other professionals  
10 working with emotionally distressed patients. Leaving GPs to rely on their own  
11 personal resources, on informal collegiate support or ad hoc relationships with  
12 colleagues in secondary care resulted in a fragile structure which could amplify  
13 rather than ameliorate anxiety.  
14  
15  
16  
17  
18

### 19 **Anxiety related to professional performance:** *across disciplinary boundaries*

20 GPs across the board expressed frustration with access to secondary care services,  
21 reporting long delays and frequent rejection of referrals, and a lack of clarity about  
22 how the services were structured and governed. GP experiences and degrees of  
23 frustration varied with an emerging picture of problematic access to services being  
24 associated with higher levels of professional anxiety. More constructive cross-  
25 disciplinary relationships were described with CAMHS workers offering clinical  
26 updates meetings and were consultants were accessible by telephone.  
27  
28  
29  
30  
31  
32  
33

### 34 **Anxiety related to interacting with young people**

35 The early finding of anxiety and uncertainty in this area was under-pinned by the  
36 difficulties GPs talked about experiencing when communicating with young people.  
37 Neither the age nor the gender of the GP appeared to facilitate communication.  
38 Female patients were generally considered to be easier to talk with whilst young men  
39 were seen to be more challenging because of their perceived reluctance to seek help  
40 and their tendency to present late.  
41  
42  
43  
44  
45  
46  
47

48 Communication difficulties included establishing a rapport, finding the right words  
49 and tone to use and dealing with silence. An inability to read the non-verbal signs,  
50 and to translate an often terse description from the young person into a coherent  
51 picture of their internal emotional state, left many GPs either relying on the  
52 accompanying parent or closing down the consultation.  
53  
54  
55  
56  
57  
58  
59  
60



1  
2  
3 Young people were seen as a highly heterogeneous group who showed variability  
4 from one presentation to the next, and also across lines of age and gender. Knowing  
5 what was 'normal' for an individual, particularly if it was presented as a principal  
6 reason for consulting with the GP, was perceived as problematic and anxiety  
7 provoking, both for the young person and for the GP.  
8  
9  
10

### 11 **Anxiety associated with the complexity of presentations of adolescent** 12 **emotional distress** 13

14  
15  
16 GPs' accounts of their experiences described a terrain beset with pitfalls, associated  
17 with the unspoken or with complex narratives embedded in social contexts. There  
18 was a sense of unpredictability and volatility to presentations which left GPs  
19 uncertain about how much input to offer at the initial consultation. This was in  
20 contrast to the rare but grave consequences which might arise when a young person  
21 seriously attempted or completed suicide; to which many GPs referred.  
22  
23  
24  
25  
26

27 Although it was accepted that uncertainty as a feature of general practice was not  
28 restricted to the clinical area of youth mental health, the early analysis showed a  
29 distinct narrative emerging in which adolescent mental health was seen as more  
30 notably anxiety provoking because of its more nebulous presentation and multiple  
31 confounding factors, largely pertaining to the social environment. The account given  
32 in the consulting room was described as the 'iceberg' indicating that often much is  
33 left hidden, or unsaid, but which nevertheless has to be raised at some point if the  
34 young person's distress is to be addressed.  
35  
36  
37  
38  
39  
40

41 Not only is there a dominant narrative of anxiety and uncertainty surrounding how  
42 GPs make sense of adolescent emotional distress, but similar responses surround  
43 management options. Few GPs expressed any degree of confidence about how they  
44 would tackle individual presentations. A small number of those with additional roles  
45 in mental health or working with patients with substance abuse problems spoke of a  
46 more systematic approach but even established GPs with personal experience of  
47 working in 'a teen drop-in clinic' or with drug dependent patients were uncertain of  
48 their practice. A paucity of treatment options was a core finding along with a lack of  
49 clarity about what GPs might reasonably do, if supported by adequate professional  
50 development.  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

## Discussion

### Summary

Anxiety and uncertainty associated with adolescent emotional distress emerged from all GP participant accounts and from the early iterative analysis of the data. Anxiety was associated with the clinical consultation, with what was expected of the GP, and how they might best respond in the absence of few clinical guidelines and limited options to involve other health and social care professionals. Unease when communicating with young people and of interpreting their accounts of distress inhibited GPs and was compounded by the complexity of presentations which ranged from familial discord to school refusal to offending behaviour. The heterogeneity of adolescent behaviour taxed GPs as did the unpredictability of the unfolding clinical presentation which could settle spontaneously or develop into a serious mental health disorder.

Whilst there was a spectrum of levels of anxiety experienced by GPs, there was a prevailing universality about the experience. How GPs responded and managed the perceived threat to professional competence and confidence was interrogated in the next stage of the analysis.

### *Strengths and limitations*

The management of adolescent mental health problems remains an under-investigated area of clinical practice. Previous studies have often been conducted by psychiatrists and whilst plurality of perspectives is important, unless more is known and understood about how GPs perceive the area many assumptions will go unchallenged. Using grounded theory, augmented by situational analysis, permits a rich exploration of the territory and facilitates theory building.

Theoretical sampling supports theory development whilst not purporting to provide universal generalizability. After 19 in-depth interviews, buttressed by situational analysis, no new themes emerged and theoretical saturation was reached. All of the respondents were white British and whilst they were recruited on the basis of their contribution to the study, it must be acknowledged that the absence of including the

1  
2  
3 experiences of GPs raised and educated in different cultural contexts will lead to the  
4 silencing of other cultural perspectives.  
5  
6

7  
8 The lead researcher and interviewer is a GP (JR). Interviewing peers has been  
9 described <sup>29</sup>as enriching the data collection because of the shared knowledge and  
10 familiarity with the clinical territory but it can lead to collusion between interviewer  
11 and respondent which needs attention and reflexivity. Co-contributors AC and JF  
12 have academic expertise in social policy and sociology which strengthened the  
13 analysis.  
14  
15  
16

### 17 18 19 *Comparison with existing literature*

20  
21 Heath asserts that *a commitment to uncertainty is fundamental to general practice:*  
22 <sup>30</sup>: 651) and Schon has described this operative landscape as a 'swampy lowland'  
23 proposing a model which advocates reflective practice as the key to dealing with  
24 uncertainty.<sup>31</sup> A quest for certainty in areas of complex practice, especially when it  
25 concerns individual experiences can be counter-productive and scholars have  
26 cautioned against clinging to the 'shelter of diagnosis' <sup>32</sup> when what is required  
27 involves attention to alleviating suffering and working purposefully with patients to  
28 catalyse their own creative capacity.<sup>15</sup> Illife et al's earlier cited work demonstrated  
29 that when GPs were fixed on the concept of depression as disease they were  
30 uncomfortable talking to young people. <sup>33 34</sup>  
31  
32  
33  
34  
35  
36  
37  
38

39 This study suggests that it is the anxiety and threat to professional competence,  
40 experienced at multiple levels, and amplified with regard to the complexity of  
41 adolescent presentations and perceived paucity of management options which  
42 compromises GPs' professional engagement and inhibits them from taking a more  
43 active role.  
44  
45  
46

### 47 48 49 *Implications for practice and research*

50  
51 Inadequate preparation, both at under and post-graduate level, is pivotal in  
52 sustaining the anxiety around clinical practice. Doctors need to be introduced to the  
53 developmental trajectory of adolescence and the conceptual framework which  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 locates adolescence as the foundation of future health <sup>35</sup> both in undergraduate  
4 education and revisited in continuing professional development .  
5  
6

7 In addition, the links between general practice and CAMHS need to be strengthened  
8 both in terms of education and understanding more of how each discipline operates,  
9 but also at a pragmatic, operational level. If cross-disciplinary practice was  
10 facilitated more treatment options would be presented at a primary care or early  
11 intervention level . Evidence of effective, feasible, primary care based brief  
12 behavioural interventions would equip GPs to engage with young people with greater  
13 confidence.  
14  
15  
16  
17  
18

19 The needs of young people are ill- served by the current provision <sup>18 19</sup> and whilst  
20 rhetoric has called for GPs to be more involved, unless we address the disabling  
21 anxiety and uncertainty in this area practice will remain static with GPs reluctant to  
22 become involved in youth mental health.  
23  
24  
25  
26

27  
28 **How this fits in**

29 GPs are known to have difficulty recognizing and responding to  
30 adolescent emotional distress. Reluctance to medicalize distress  
31 has been reported.  
32

33 This study shows that anxiety and uncertainty about practice in this  
34 complex clinical area are universal and independent of age, gender,  
35 level of experience of GP.  
36  
37

38 If GPs are to play a more active role in the early identification and  
39 intervention of distress we need to know more about the factors  
40 which ameliorate or exacerbate professional anxiety about  
41 practice. Critically, adolescent mental health needs to feature in  
42 undergraduate and postgraduate curricula.  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

Table 1

Participant Number	Gender	Age	Salaried or Partner	Practice descriptor	Additional professional experience
01	F	50-59	S	Semi-rural Deprived	GP Postgraduate education
02	Male	50-59	S	Urban Deprived	Addiction medicine in primary care
03	Female	50-59	P	Urban Deprived; wealthy student population	Former Assoc. Specialist in CAMHS
04	Female	40-49	S	Semi-rural Deprived	Mental health Lead for a PCT
05	Female	20-29	S	Urban Deprived	
06	Male	40-49	P	Semi-rural Largely affluent	
07	Male	40-49	P	Semi-rural Mixed :	Child Protection Lead for a PCT
8	Female	30-39	S	Semi-rural Mixed :	
9	Male	50-59	P	Semi-rural Mixed :	GP lead for 'teen drop-in' clinic
10	Male	40-49	P	Urban Deprived	Mental Health and Child Protection Lead for a PCT. Substance misuse

11	Female	20-29	S	Urban	
				Deprived	
12	Male	30-39	S	Semi-rural	
				Mixed: largely affluent	
13	Female	30-39	S	Urban	
				Deprived	
14	Male	40-49	P	Urban	
				Deprived	
15	Male	40-49	P	Semi-rural	
				Mixed :	
16	Female	20-29	S	Urban	
				Deprived	
17	Male	30-39	S	Urban	
				Deprived	
18	Female	40-49	P	Semi-rural	
				Affluent	
19	Female	50-59	P	Semi-rural	Child health lead
				Mixed :	

## Anxiety paper Boxes i

Box 1. Anxiety related to professional performance: In the consultation

*I've always thought young people are challenging and still do and I have more questions than answers' (06;M; 40-49;P)*

*so not knowing what to do is a bit of a theme really (07;M; 40-49;P)*

*I find the adults will accept me at face value, generally. And they come with something usually fairly clear and they want that sorting out, it might not be straight forward, it might not even be simple they might have even brought things off the internet but it is a fairly clear baggage package... what I find with younger people with psychological or emotional disorders is it's not a clearly packed problem, it's in the extreme realms of the undefined. (06;M;40-49;P)*

*I know we don't meet all the people we refer to but we usually have something to hang it on because of our experience as junior doctors working in hospital but with CAMHS there is nothing. (08; F;30-39;S)*

Box 2 Anxiety related to professional performance: at a structural level

*NICE guidelines a few years ago looked at depression in young people and kind of hampered our ability to do anything with them really (07;M; 40-49;P)*

*'...because it doesn't fit within any ticky box guidelines until time has passed I rarely know whether I've done the right thing, it's all in retrospect. (06;M;40- 49;P)*

*I'll bring people back in 1 week, I don't think this annoys my partners but it can become a bugbear....I'll squeeze them in when there are no appointments, which is probably making a rod for my own back and I wouldn't encourage trainees to do it, but I like the idea of seeing something through to its natural conclusion...its perhaps my own insecurity (14;M; 40-49,P)*

*What we don't have, in general practice is supervision...no counsellors are allowed to work without it but GPs are just sent out there, andI really do feel there is a huge need for it even if it is just one phone call-it's that ability to share the responsibility, not to dump it, but to genuinely share it.(04;F; 40-49;S)*

## Anxiety Boxes ii

Box 3. Anxiety related to professional performance: across disciplinary boundaries

*Some (mental health) creatures are on the verge of being mythical beasts, ...like psychotherapists,...educational psychologists (09;M;50-59;P)*

*CAMHs..it feels like it's a bit of a hotchpotch really, a patched together sort of service and I'm not sure who is control....people who counsel children-I don't know much about them, how much responsibility they take.. (07;M; 4049;M)*

Box 4. Anxiety related to interacting with young people

*Generally consulting with young people, I often find, if I'm being honest, probably more difficult than I would expect to find it. I think I probably have this unrealistic view of myself as really sort of approachable and you know still being quite young myself compared to other GPs, being able to communicate fairly easily and fairly well with young people, then always very quickly, it becomes apparent that actually no, you are a million miles away from where they are, and they don't really relate to you very well at all....(08; F; 30-39; S)*

*Males with mental health issues worry me intensely, it really does seem that there is not an awful lot of trivia goes on there. By the time a male is presenting, because they don't have the tools to come to the GP very often, they don't understand that you can just come along when things are in their development, they usually come when something is really big ,black and bleak.(14;M;40-49;P)*

*I struggle a bit to work out how to word sort of mental health questions with to the sort of under 16 year olds particularly.... I suppose with adults I have my kind of, standard questions ..but using those sort of questions with young people often draws a blank face, and, so it's something I have to rephrase; I feel that I don't necessarily know their kind of lingo if you like...(17;M; 30-39;S)*

*So he went off to do a urine sample and I was pleased to speak to his parents without him, seemed easier to talk about some of the mental issues without him there... (017;M; 30-39;S)*

*With children and teenagers it tends to be you controlling the pace of the consultation.... and you finish the consultation when you want to (07;M; 40-49;P)*



## Anxiety Box iii

Box 4. Anxiety related to interacting with young people

*there's no such thing as a typical teenage presentation ( 13;F; 30-29;S)*

*...and her mum said, as mum's do, oh she has dreadful mood swings on her periods, it's like living with a monster ..and then they say 'but that's normal isn't it, she is 14?'... so it's hard to know what to do (11;F; 20-29;F)*

Box 5. Anxiety associated with the complexity of presentations of adolescent emotional distress

*... you feel that there are these big 'no go areas' in teenage consultations which loom over you like a black cloud (11;F; 20-29; S)*

*They are missing school, in trouble with the police, youth offending team or not uncommonly the parent just comes by themselves...I never get a 14 year old acting out saying 'you know, I'm in trouble with my mam and dad' ( 10;M;40-49;P)*

*they are in absolute crisis one minute and then you see them a week later and they can hardly remember what it was all about (07;M; 40-49;P)*

*Its always a worry isn't it that you just completely get it wrong..I mean I'm conscious of this. I had someone in on Monday, parents, whose son had just hung himself at 21. I'd never seen him, he was a patient here. They had no idea anything was wrong. Nobody did....there is always that underlying things isn't there, that you might miss something catastrophic.. (01;F;50-59;S)*

## Anxiety Box iii

Box 5. Anxiety related to interacting with young people

*there's no such thing as a typical teenage presentation ( 13;F; 30-29;S)*

*...and her mum said, as mum's do, oh she has dreadful mood swings on her periods, it's like living with a monster ..and then they say 'but that's normal isn't it, she is 14?'... so it's hard to know what to do (11;F; 20-29;F)*

Box 6. Anxiety associated with the complexity of presentations of adolescent emotional distress

*... you feel that there are these big 'no go areas' in teenage consultations which loom over you like a black cloud (11;F; 20-29; S)*

*They are missing school, in trouble with the police, youth offending team or not uncommonly the parent just comes by themselves...I never get a 14 year old acting out saying 'you know, I'm in trouble with my mam and dad' ( 10;M;40-49;P)*

*they are in absolute crisis one minute and then you see them a week later and they can hardly remember what it was all about (07;M; 40-49;P)*

*Its always a worry isn't it that you just completely get it wrong..I mean I'm conscious of this. I had someone in on Monday, parents, whose son had just hung himself at 21. I'd never seen him, he was a patient here. They had no idea anything was wrong. Nobody did....there is always that underlying things isn't there, that you might miss something catastrophic.. (01;F;50-59;S)*

*Uncertainty is very key to this group when you're looking - in terms of depression and suicide risk and things like that, you know, it's standard. Young people particularly young males are quite at risk of just going off and doing something. (04; F; 40-49;S)*

*The main anxiety is what to do. (07;M; 40-49;P)*

1  
2  
3 **Acknowledgments.** The authors wish to thank the GP participants for their time and  
4 candour, and their practises for supporting the research; and the RCGP Scientific  
5 Foundation Board which awarded a grant to cover the transcription costs.  
6

7 **Competing interests.** We declare no competing interests.  
8

9  
10 **Funding:** the RCGP Scientific Board awarded a grant to cover the transcription  
11 costs. SFB-2008-06.  
12

13 **Ethical approval:** Hull & East Riding Local Ethics Committee. REC Reference No:  
14 08/H1304/97.  
15

16 **Data sharing:** Extra data is available by emailing Jane Roberts on  
17 jane.roberts@sunderland.ac.uk  
18

19  
20 This includes theoretical memos, field notes, anonymized transcripts and situational  
21 analysis diagrams.  
22

23 **Contributorship statement:** JR was the lead investigator , conducted all of the  
24 interviews, carried out the primary analysis of the data and wrote the manuscript  
25 AC and JF contributed to the design of the study and met regularly with JR to look at  
26 the data and the analysis at each stage to agree on the open, axial and selective  
27 codes  
28

29  
30 AC was involved with GP recruitment and commented on each draft of the  
31 manuscript including the final submission  
32

33 JF read all drafts of the manuscripts and agreed with AC's final comments  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

## Reference list

1. Green H, McGinnity A, Meltzer H, Ford T, Goodman R. Mental Health of Children and Young People in Great Britain, 2004 : summary report. London: Office of National Statistics, 2005.
2. Singleton N, Bumpstead R, O'Brien M, Lee A, Meltzer H. Psychiatric morbidity in adults living in private households, 2000. Norwich: The Stationery Office, 2001.
3. Hawton K, Saunders KE, O'Connor RC. Self-harm and suicide in adolescents. *Lancet* 2012;379(9834):2373-82.
4. Kramer T, Garralda M. Child and adolescent mental health problems in primary care. *Advances in Psychiatric Treatment* 2000;6:287-94.
5. Hickie I, Fogarty A, Davenport T, Luscombe G, Burns J. Responding to experiences of young people with common mental health problems attending Australian general practice. *Medical Journal of Australia* 2007;187:S47-S52.
6. Gledhill J, Garralda M. The short-term outcome of depressive disorder in adolescents attending primary care: a cohort study. *Soc Psychiatry Psychiatr Epidemiol* 2011;46(10):993-1002.
7. Martinez R, Reynolds S, Howe A. Factors that influence the detection of psychological problems in adolescents attending general practices. *Br J Gen Pract* 2006;56(529):594-9.
8. Haller D, Sanci L, Sawyer S, Patton G. The identification of young people's emotional distress: a study in primary care. *Br J Gen Pract* 2009;59(560):e61-70.
9. Mauerhofer A, Berchtold A, Michaud PA, Suris JC. GPs' role in the detection of psychological problems of young people: a population-based study. *Br J Gen Pract* 2009;59(566):e308-14.
10. Churchill R. Child and Adolescent Mental Health. In: Cohen A, editor. *Delivering mental health for primary care: an evidence based approach*. London: RCGP, 2008:157-84.
11. Iliffe S, Gledhill J, da Cunha F, Kramer T, Garralda E. The recognition of adolescent depression in general practice: issues in the acquisition of new skills. *Primary Care Psychiatry* 2004;9(2):51-56.
12. Goldberg D, Huxley P. *Common mental disorders: a bio-social model*. London: Tavistock/Routledge, 1992.
13. Pilling S, Whittington C, Taylor C, Kendrick T. Identification and care pathways for common mental health disorders: summary of NICE guidance. *BMJ* 2011;342:d2868.
14. Dowrick C. *Beyond Depression: a new approach to understanding and management*. 2nd ed. Oxford: Oxford University Press, 2009.
15. Reeve J. Interpretive medicine: Supporting generalism in a changing primary care world. *Occas Pap R Coll Gen Pract* 2010(88):1-20, v.
16. Dowrick C, May C, Richardson M, Bundred P. The biopsychosocial model of general practice: rhetoric or reality? *Br J Gen Pract* 1996;46(403):105-7.
17. Karasz A, Dowrick C, Byng R, Buszewicz M, Ferri L, Hartman TCO, et al. What we talk about when we talk about depression: doctor-patient conversations and treatment decision outcomes. *Br J Gen Pract* 2012;62(594):e55-e63.
18. Kennedy I. Getting it right for children and young people: overcoming cultural barriers in the NHS so as to meet their needs. A review by Sir Ian Kennedy. London: Department of Health, 2010.
19. Layard R. How mental health loses out in the NHS. London: London School of Economics and Political Science, 2012.
20. Department for Education and Skills. Every Child Matters: change for children London: HMSO, 2002.
21. Department of Health. *Child and Adolescent Mental Health-National Service Framework for Children, Young People and Maternity Services* London: Department of Health 2004.
22. Department of Health. *Healthy Lives, Brighter Futures* London: The Stationery Office, 2009.

23. National Institute for Health & Clinical Excellence. Depression in children and young people: identification and management in primary, community and secondary care. *National Clinical Practice Guidelines*. London: NICE, 2005.
24. Tait L. To disclose or not to disclose psychological problems to GPs. *Br J Gen Pract* 2009;59(566):638-9.
25. Biddle L, Donovan JL, Gunnell D, Sharp D. Young adults' perceptions of GPs as a help source for mental distress: a qualitative study. *Br J Gen Pract* 2006;56(533):924-31.
26. Glaser B, Strauss A. *The discovery of grounded theory*. Chicago: Aldine, 1967.
27. Charmaz K. *Constructing Grounded Theory: A practical guide through qualitative analysis*. Thousand Oaks, California: Sage Publications, Inc, 2006.
28. Clarke A. *Situational analysis: Grounded Theory after the postmodern turn*. Sage Publications, Inc. Thousand Oaks, California: Sage Publications, Inc, 2005.
29. Chew-Graham CA, May CR, Perry MS. Qualitative research and the problem of judgement: lessons from interviewing fellow professionals. *Family Practice* 2002;19(3):285-89.
30. Heath I. Uncertain clarity: contradiction, meaning, and hope. *Brit J Gen Pract* 1999;49(445):651-57.
31. Schon D. *The reflective practitioner*. San Francisco: Jossey-Bass, 1983.
32. Dowrick C. Reasons to be cheerful ? Reflections on GPs' responses to depression. *Br J Gen Pract* 2009;59(566):636-7.
33. Iliffe S, Williams G, Fernandez V, Vila M, Kramer T, Gledhill J, et al. Treading a fine line: is diagnosing depression in young people just medicalising moodiness? *Br J Gen Pract* 2009;59(560):156-7.
34. Iliffe S, Gallant C, Kramer T, Gledhill J, Bye A, Fernandez V, et al. Therapeutic identification of depression in young people: lessons from the introduction of a new technique in general practice. *Br J Gen Pract* 2012;62(596):e174-82.
35. Sawyer SM, Afifi RA, Bearinger LH, Blakemore SJ, Dick B, Ezech AC, et al. Adolescence: a foundation for future health. *Lancet* 2012;379(9826):1630-40.

## Appendix 1 Early Topic Guide

1. I'd like to talk about your experiences of consulting with young people in general
  - How do you find this age group?
  - Is it very different to consulting with older patients?
  - What sort of problems do you see? Do they consult often?
2. Can we talk more about consulting with young people who may have psychological/mental health problems
  - How do you find this clinical area?
  - What about seeing YP alone/ with 'another'
  - Any areas particularly tricky to broach ?
3. How do you consider possible 'mental health problems' which presenting in young people ?
  - Do any examples come to mind ?
  - What approach did you take
  - What worked well? What was difficult?
  - Is it different with other age groups
4. What are your thoughts on 'depression' and 'anxiety' in young people ?
  - Do you see much of it?
  - Does this differ from other age groups?
  - What options are there in primary care?
5. Do you think GPs have a role/or not in promoting emotional well-being in young people? Explore

1  
2  
3  
4  
5 Research checklist  
6  
7  
8

9 As this is a qualitative study it does not fall within the parameters of the recommended research  
10 checklists.  
11

12  
13  
14 A statement to this effect is included in the covering letter.  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60



**“.. I think this is maybe our Achilles Heel....” Exploring GPs’ responses to young people presenting with emotional distress in general practice. A qualitative study.**

Journal:	<i>BMJ Open</i>
Manuscript ID:	bmjopen-2013-002927.R1
Article Type:	Research
Date Submitted by the Author:	10-Jul-2013
Complete List of Authors:	Roberts, Jane; University of Sunderland, Pharmacy, Health & Well-being Crosland, Ann; University of Sunderland, Pharmacy, Health & Well-being Fulton, John; University of Sunderland, Pharmacy, Health & Well-being
<b>Primary Subject Heading</b>:	General practice / Family practice
Secondary Subject Heading:	Mental health, Medical education and training, Qualitative research
Keywords:	PRIMARY CARE, MENTAL HEALTH, Child & adolescent psychiatry < PSYCHIATRY

SCHOLARONE™  
Manuscripts



1  
2  
3  
4  
5 **“.. I think this is maybe our Achilles Heel....” Exploring GPs’ responses to**  
6 **young people presenting with emotional distress in general practice. A**  
7 **qualitative study.**  
8  
9

10  
11  
12  
13  
14 Jane H Roberts , Ann Crosland, John Fulton

15  
16 Department of Pharmacy, Health & Well-being, Faculty of Applied Sciences, The  
17 Science Complex, Wharnccliffe St, Sunderland, SR1 3SD.

18  
19  
20 Jane H Roberts, Clinical Senior Lecturer, University of Sunderland,

21  
22 Ann Crosland, Professor of Nursing , University of Sunderland,

23  
24 John Fulton, Principal Lecturer, University of Sunderland,  
25  
26  
27

28  
29 Correspondence to Jane H Roberts

30  
31 jane.roberts@sunderland.ac.uk  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 **“.. I think this is maybe our Achilles Heel....” Exploring GPs’ responses to**  
4 **young people presenting with emotional distress in general practice. A**  
5 **qualitative study.**  
6  
7  
8  
9

## 10 **Article summary**

### 11 **Article Focus:**

- 12 1. an exploratory study
- 13 2. to examine GPs’ views and experiences of consulting with young people
- 14 3. to better understand GPs perspectives

### 15 **Key Messages**

- 16 1. Anxiety about practice experienced when consulting with young  
17 people is the dominant finding in a first stage analysis of a qualitative study.  
18 This is independent of age and gender of GP
- 19 2. Anxiety relates to professional performance; interacting with young  
20 people and the complex nature of presentations of emotional distress in  
21 primary care
- 22 3. Unless anxiety and related uncertainties about practice are  
23 addressed GPs will continue to miss opportunities to address early  
24 emotional difficulties and young people’s mental health needs in primary care  
25 will continue to be poorly met

### 26 **Strengths and Limitations**

- 27 1. Qualitative research in under -examined areas offers new insights  
28 and explores why behaviours might arise
- 29 2. The data presented contributes to theory building
- 30 3. Theoretical sampling led to only white British born GPs participating  
31 so other cultural perspectives were not included

1  
2  
3 **“.. I think this is maybe our Achilles Heel....” Exploring GPs’ responses to**  
4 **young people presenting with emotional distress in general practice. A**  
5 **qualitative study.**  
6  
7  
8  
9

10  
11 **Main text**  
12

13 **Introduction**  
14

15  
16 Emotional distress in young people is common. It may be the affective response to  
17 the challenges of everyday life or may indicate a mental health disorder compatible  
18 with a psychiatric diagnosis. The most recent and widely cited household survey  
19 reports at least 10% of 10-15 year olds <sup>1</sup> and 17 % of 16-19 year olds <sup>2</sup> to ) to have  
20 symptoms consistent with a mental health disorder as defined by the ICD-10.  
21 Behavioural manifestations of emotional distress might include self-harm which, at a  
22 conservative estimate, appears to affect around 10% of adolescents, as reported in  
23 six studies cited by Hawton et al in a recently published review . <sup>3</sup>  
24  
25  
26  
27  
28  
29

30  
31 Data from populations of young people who consult their GP reveal higher rates of  
32 psychological distress, of the order of 20-30%. <sup>4 5</sup> GPs identify serious mental illness  
33 but often fail to detect less severe manifestations <sup>6</sup> and appear reluctant to discuss  
34 emotional issues<sup>7</sup>; unless offered cues by the young person in the consultation <sup>8</sup> or if  
35 other factors are present such as a previous history of a suicide attempt or a pattern  
36 of frequent consulting <sup>9</sup>. Young people’s presentations in primary care are often  
37 complex and present with behavioural, psychosocial, academic and familial  
38 problems which can be problematic to untangle in contrast to adult mental health  
39 manifestations which, although variable, may be less intense in their presentation .  
40 Adolescent emotional distress may indicate underlying co-morbid mental health  
41 problems and it It has been suggested that often the ‘most important features in  
42 terms of assessment may be concealed or hidden’. <sup>10</sup>. A key concern is the difficulty  
43 of distinguishing between ‘moodiness’ or a persisting emotional disorder and GPs  
44 have expressed a worry at ‘over-medicalising young people’s lives’. <sup>11</sup> Illiffe &  
45 colleagues found that GPs were uncomfortable about making a diagnosis of  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 depression in young people (the most common, but often coexisting, mental health  
4 problem in adolescence ).  
5  
6

7 On the other hand GPs' are increasingly involved in managing common mental  
8 health problems in older patients <sup>12</sup> Although a biomedical perspective dominates,  
9 supported by an array of NICE clinical guidelines, alternative frameworks for  
10 considering adult mental health problems have been offered . Dowrick <sup>13</sup> and Reeve  
11 <sup>14</sup> refer to the insights derived from the wisdom traditions in informing their work  
12 which moves away from a positivist understanding of emotional distress to an  
13 approach which incorporates ideas of personal agency and encourages hope. <sup>15</sup>  
14 Historically, research has found GPs to be largely dismissive of their role in  
15 addressing social issues in adult mental ill-health <sup>16</sup> . Contemporary studies reveal a  
16 shift with greater awareness of the lay perspective, which typically favours a social  
17 model adult mental ill-health <sup>17</sup> , and a matched response by GPs mirroring popular  
18 social constructions of distress .  
19  
20  
21  
22  
23  
24  
25  
26  
27

28 Despite the challenge of responding to emotional distress in adolescence and the  
29 patchy, often inadequate provision of secondary care services <sup>18 19</sup> a series of policy  
30 directives have emphasised the role of GPs and other front-line services, in the  
31 promotion of psychological well-being and the early indication of difficulties. <sup>20 21 22</sup>  
32 Practitioners are expected to have 'sufficient knowledge, training and support 'in this  
33 area including competence in 'active listening' and conversational technique' <sup>23</sup> .  
34  
35  
36  
37  
38

39 There is a growing body of evidence examining young people's experiences of  
40 talking to GPs about emotional problems. They reveal a mixed picture including a  
41 reluctance to disclose <sup>24</sup> , a fear of being judged or offered medication <sup>25</sup> . Much less  
42 is known about GP perspectives. This paper presents a qualitative, exploratory study  
43 which examines GPs' views and experiences of consulting with young people  
44 presenting with emotional distress.  
45  
46  
47  
48

## 49 **Method**

### 50 **Study Design**

51  
52  
53  
54 The study took place in the North East of England in 18 general practices based in  
55 urban, rural and semi-rural communities serving predominantly socio-economically  
56  
57  
58  
59  
60

1  
2  
3 disadvantaged patients. The qualitative study comprised of in depth individual  
4 interviews with GPs recruited using theoretical sampling. As early theoretical ideas  
5 emerged successive GPs were recruited on the basis of their capacity to contribute  
6 to the development or abandonment of initial theoretical constructs.  
7  
8

9  
10 Data were collected between January 2010 to May 2011  
11

### 12 *Participants*

13  
14 GPs with less than four years clinical experience were excluded. The initial recruits  
15 were selected on the basis of their relevant experience and their ability to generate  
16 early data which would scope the terrain of the area under enquiry; for example  
17 having a role as mental health lead or previous experience working in Child &  
18 Adolescent Mental Health services (CAMHS)  
19  
20

21 GPs were approached by telephone and email contact and sent information sheets.  
22 A follow-up contact established their verbal consent to meet at a location of their  
23 choice. Two GPs approached declined to participate. One cited forthcoming  
24 extended annual leave and another a view that as the senior partner he saw  
25 relatively few younger aged patients and suggested recruitment of a younger GP in  
26 the same practice.  
27  
28

29 Ethical approval by the Local Research Ethics Committee, the seven Primary Care  
30 Trust organizations of the region and the University of Sunderland was granted  
31 before data collection began.  
32  
33

### 34 *Data collection and analysis*

35 The audio-taped semi-structured interviews were transcribed verbatim with consent.  
36 An initial topic guide was used with the first tranche of participants based on the  
37 extant literature and developed through discussion. The topic guide was then revised  
38 on the basis of ideas arising from the early interviews, and the iterative analysis  
39 which began as soon as the first interview was undertaken. The interview guides  
40 explored doctors' experiences of consulting with young people in general and those  
41 presenting with psychological or mental health problems, GPs' understanding of  
42 depression and anxiety in adolescence, of how emotional distress presents in the  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 surgery and the role of the GP in promoting emotional well-being in young people  
4 (See appendix 1). The guide was refined to include questions about how structural  
5 changes impacted on, and consultation style shaped, practice.  
6  
7

8  
9 The interviews lasted between 50 to 75 minutes. Field notes and theoretical memos  
10 were kept throughout the period of data collection and analysis.  
11

12  
13 The transcripts were coded and analysed using the grounded theory method  
14 described by Strauss and Glaser<sup>26</sup> and revised by Charmaz.<sup>27</sup> The constant  
15 comparative method of analysis is core to the process and informs the theoretical  
16 sampling of recruits. Early ideas were tested with subsequent participants and found  
17 to be either substantiated or rejected through the iterative process of constant  
18 comparison supported by theoretical sampling. Situational maps, both 'messy' and  
19 'ordered', were constructed during this phase of analysis.<sup>28</sup>  
20  
21  
22  
23

24  
25 The data presented here were generated after the first level of analysis was  
26 completed, during which only the open codes were iteratively developed by JR and  
27 subjected to further examination by AC (primary care academic) and JF(sociologist).  
28 Further analysis of the axial and selective codes will be presented in two subsequent  
29 companion papers.  
30  
31  
32  
33

### 34 **Results**

35  
36  
37 Nineteen GPs participated, 10 women. (Table1). The early iterative analysis of the  
38 data found the open codes to support a dominant narrative of anxiety under-pinning  
39 the majority of the research interviews. This pervasive and disabling emotional  
40 response to encounters with emotionally distressed young people appeared to  
41 coalesce around three domains.  
42  
43  
44

45  
46 These can be viewed as anxiety experienced by GPs in response to:  
47

- 48 1) professional performance; in the consultation, at an external level, across  
49 disciplinary boundaries;  
50
- 51 2) interacting with young people; and  
52
- 53 3) the complexity of presentations of adolescent emotional distress  
54  
55  
56  
57  
58  
59  
60

Each of the three themes will be presented in turn and supported by illustrative quotations taken from the transcripts (see boxes 1-6). GP participants are identified by identifier number, gender, age range and whether salaried or a partner ( as presented in Table 1.)

**1. i. Anxiety related to professional performance:** operating *In the consultation*

A coherent narrative emerged, gathered from almost all of the participants, of practitioners being anxious in the consultation because of an uncertainty about what to do and of what was expected of them, as primary care clinicians.

This resulted in a sense of professional impotence. It was acknowledged that feeling uncertain about how best to proceed, and unsure of practice, led to a sense of disempowerment through not knowing what to do. This was in contrast to accounts of working with older patients where the options for GPs appear more clearly defined. The data generated by the open code analysis suggested that not being able to formulate the initial presentation by a young person into a definable 'disorder' created a sense of operating in uncharted territory.

Anxiety was amplified by the lack of exposure to adolescent mental health in undergraduate medical education which was the unanimous experience of all participants. Where the topic had been included in the curriculum, it was often restricted to severe mental disorder for example being assigned to medical teams looking after adolescents hospitalized with anorexia nervosa.

(See box 1).

**1. ii Anxiety related to professional performance:** operating *at an external level*

A lack of benchmarks in practice meant assessing one's performance in relation to peers was problematic since no 'gold standard' existed. The only NICE guideline which was referenced (concerning the management of depression in under 18 year olds) was regarded as having "hampered GPs" from becoming involved in the management of adolescent depression since the Guideline did not advocate the use



1  
2  
3 of anti-depressants and, with access to psychological therapies piecemeal, appeared  
4 to support a position that there was little to be offered in primary care.  
5  
6

7 Constraints in practice led to frustration and an anxiety about management. For  
8 example, varying arrangements within practices governing access to appointments  
9 and the ease, or not, of maintaining continuity of care were seen to contribute to  
10 professional anxiety by impeding attentive 'watchful waiting' and some GPs  
11 described attempts to circumvent inflexible appointment systems in order to be more  
12 available to patients.  
13  
14  
15  
16

17 A lack of professional supervision was identified by a small number of more  
18 experienced GPs involved with Postgraduate Training and provision of mental  
19 health services at a regional level, and contrasted to systems for other professionals  
20 working with emotionally distressed patients. Leaving GPs to rely on their own  
21 personal resources, on informal collegiate support or ad hoc relationships with  
22 colleagues in secondary care resulted in a fragile structure which could amplify  
23 rather than ameliorate anxiety.  
24  
25  
26  
27  
28

29  
30 (See box 2).  
31

32  
33 **1. iii Anxiety related to professional performance: across disciplinary**  
34 *boundaries*  
35  
36

37 GPs across the board expressed frustration with access to secondary care services,  
38 reporting long delays and frequent rejection of referrals, and a lack of clarity about  
39 how the services were structured and governed. GP experiences and degrees of  
40 frustration varied with an emerging picture of problematic access to services being  
41 associated with higher levels of professional anxiety. Where GPs described more  
42 constructive cross-disciplinary relationships, with CAMHS practitioners offering  
43 clinical updates meetings, and where consultants were accessible by telephone, less  
44 anxiety was voiced.  
45  
46  
47  
48  
49

50 Unfamiliarity with the roles and responsibilities of CAMHS practitioners, coupled with  
51 an obligation to refer in the absence of other options, left some GPs feeling uncertain  
52 about the clinical care pathway and unsure about practice.  
53  
54  
55  
56  
57  
58  
59  
60



(See box 3.)

## 2. Anxiety related to interacting with young people

The open codes showed a dominant finding of GPs expressing anxiety associated with difficulties experienced when communicating with young people in general. Neither the age nor the gender of the GP appeared to facilitate communication, with younger and female GPs similarly as uneasy as older male and female GPs. Female patients were generally considered to be easier to talk with whilst young men were seen to be more challenging because of their perceived reluctance to seek help and their tendency to present late.

Communication difficulties included establishing a rapport, finding the right words and tone to use and dealing with silence. An inability to read the non-verbal signs, and to translate an often terse description from the young person into a coherent picture of their internal emotional state, left many GPs either relying on the accompanying parent or closing down the consultation. Being able to find common ground was identified as being key to beginning the process of establishing rapport.

Young people were seen as a highly heterogeneous group who showed variability from one presentation to the next (intra-variability), and also across lines of age and gender (inter-variability). Knowing what was 'normal' for an individual, particularly if it was presented as a principal reason for consulting with the GP, was perceived as problematic and anxiety provoking, both for the young person and for the GP.

(See box 4)

## 3. Anxiety associated with the complexity of presentations of adolescent emotional distress

GPs' accounts of their experiences consulting with young people experiencing distress described a terrain beset with pitfalls, associated with the unspoken or with complex narratives embedded in social contexts; and presented in an undifferentiated form. GPs spoke of a sense of unpredictability and volatility to presentations which left them uncertain about how the patient narrative might unfold and how much input to offer at the initial consultation. In particular this generated

1  
2  
3 anxiety associated with the rare but grave consequences which might arise when a  
4 young person seriously attempted or completed suicide; a clinical experience to  
5 which many GPs referred and which could lead to enduring professional anxiety.  
6  
7 (See box 5).  
8  
9

10 Although it was accepted that uncertainty as a feature of general practice was not  
11 restricted to the clinical area of youth mental health, the first stage analysis showed  
12 a distinct narrative emerging in which adolescent mental health was seen as more  
13 notably anxiety provoking because of its more nebulous presentation and multiple  
14 confounding factors, which largely pertained to the social environment. The account  
15 given in the consulting room was described as the 'iceberg' indicating that often  
16 much is left hidden, or unsaid, but which nevertheless has to be raised at some point  
17 if the young person's distress is to be addressed.  
18  
19

20 Not only is there a dominant narrative of anxiety surrounding how GPs make sense  
21 of adolescent emotional distress, but similar responses are associated with  
22 management options. Few GPs expressed any degree of confidence about how they  
23 would tackle individual presentations. A small number of those with additional roles  
24 in mental health or working with patients with substance abuse problems spoke of a  
25 more systematic approach to organizing and offering care. However even  
26 established GPs with personal experience of working in 'a teen drop-in clinic' or with  
27 drug dependent patients described uncertainty about their practice. A paucity of  
28 treatment options was a consistent finding along with a lack of clarity about what  
29 GPs might reasonably be expected to do, if supported by adequate professional  
30 development.  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42

## 43 Discussion

### 44 Summary

45  
46  
47 Anxiety about practice, coupled with a perceived reduced range of options and lack  
48 of clarity of expectations, associated with diverse presentations of adolescent  
49 emotional distress in primary care, emerged from all GP participant accounts in the  
50 first stage of analysis. Unease when communicating with young people and  
51 difficulties interpreting their accounts of distress inhibited GPs. This was  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 compounded by the complexity of presentations which ranged from familial discord  
4 to school refusal to offending behaviour, usually in the absence of any clear  
5 diagnosis. The heterogeneity of adolescent behaviour taxed GPs as did the  
6 unpredictability of the unfolding clinical presentation which might settle  
7 spontaneously or might develop into a serious mental health disorder.  
8  
9

10  
11  
12 Whilst there was a spectrum of levels of anxiety experienced by GPs, there was a  
13 prevailing universality about the experience. How GPs responded and managed the  
14 perceived threat to professional competence and confidence was interrogated in the  
15 next stage of the analysis which would lead to the development of the axial codes ,  
16 or pillars, of the emerging conceptual model (presented elsewhere).  
17  
18  
19

### 20 21 22 *Strengths and limitations*

23  
24 The management of adolescent mental health problems remains an under-  
25 investigated area of clinical practice. Previous research has largely been conducted  
26 by psychiatrists whose perspective is different to that of GPs responding to  
27 undifferentiated distress in the consulting room. Using grounded theory, augmented  
28 by situational analysis, permits a rich exploration of the territory and facilitates theory  
29 building.  
30  
31  
32  
33  
34

35  
36 Theoretical sampling supports theory development whilst not purporting to provide  
37 universal generalizability. After 19 in-depth interviews, buttressed by situational  
38 analysis, no new themes emerged and theoretical saturation was reached. All of the  
39 respondents were white British and whilst they were recruited on the basis of their  
40 contribution to the study, it must be acknowledged that the absence of including the  
41 experiences of GPs raised and educated in different cultural contexts will lead to the  
42 silencing of other cultural perspectives.  
43  
44  
45  
46  
47

48  
49 The lead researcher and interviewer is a GP (JR). Interviewing peers has been  
50 described <sup>29</sup>as enriching the data collection because of the shared knowledge and  
51 familiarity with the clinical territory but it can lead to collusion between interviewer  
52 and respondent which needs attention and reflexivity. Co-contributors AC and JF  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 have academic expertise in social policy and sociology which strengthened the  
4 analysis.  
5  
6

### 7 *Comparison with existing literature*

8  
9  
10 Heath asserts that *a commitment to uncertainty is fundamental to general practice:*  
11 <sup>30</sup>: 651) and Schon has described this operative landscape as a 'swampy lowland'  
12 proposing a model which advocates reflective practice as the key to dealing with  
13 uncertainty.<sup>31</sup> A quest for certainty in areas of complex practice, especially when it  
14 concerns individual experiences can be counter-productive and scholars have  
15 cautioned against clinging to the 'shelter of diagnosis' <sup>15</sup> when what is required  
16 involves attention to alleviating suffering and working purposefully with patients to  
17 catalyse their own creative capacity.<sup>14</sup> Illife et al's earlier cited work demonstrated  
18 that when GPs were fixed on the concept of depression as disease they were  
19 uncomfortable talking to young people.<sup>32 33</sup>  
20  
21  
22  
23  
24  
25  
26  
27

28 This study which contributes to building a theoretical model, suggests that anxiety  
29 and, perceived threats to professional competence, can be experienced at multiple  
30 levels, and are amplified with regard to the complexity of adolescent presentations.  
31 This can compromise GP's professional engagement with young people.  
32 Understanding more about why some GPs can creatively respond to the anxiety and  
33 lack of certainty about expectations defines the next stage of the analysis.  
34  
35  
36  
37  
38

### 39 *Implications for practice and research*

40  
41 Inadequate educational preparation, both at under and post-graduate level, is pivotal  
42 in failing to address the anxiety around clinical practice. Doctors need to be  
43 introduced to the developmental trajectory of adolescence and the conceptual  
44 framework which locates adolescence as the foundation of future health <sup>34</sup> both in  
45 undergraduate education and revisited in continuing professional development .  
46 This approach will help GPs to understand more about why addressing emotional  
47 distress in the second decade of life is important.  
48  
49  
50  
51  
52  
53

54 GPs need good quality educational exposure and preparation to deal with the multi-  
55 axial development of adolescence and the emergence of mental health disorders in  
56  
57  
58  
59  
60

1  
2  
3 the 10 to 20 year olds. The current psychiatric classification systems do not facilitate  
4 clinical practice in this domain at primary care level. In addition, the links between  
5 general practice and CAMHS need to be strengthened both in terms of education  
6 and understanding more of how each discipline operates, but also at a pragmatic,  
7 operational level. If cross-disciplinary practice was facilitated more treatment options  
8 would be presented at a primary care or early intervention level.  
9

10  
11  
12  
13  
14 More research is needed to demonstrate evidence of effective, feasible, primary care  
15 based brief behavioural interventions which would equip GPs to engage with young  
16 people with greater confidence and support the development of evidence based  
17 policy .  
18

19  
20  
21 At a systemic level, this study shows that external factors are important in influencing  
22 practice and can moderate or exacerbate levels of anxiety. Systems which improve  
23 access to care for young people need to be introduced at practice level and  
24 supported by policy.  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38

39 The study was given ethical approval by the Hull and East Riding LREC. REC  
40 Reference No. 08/H1304/97 and the University of Sunderland's Ethics Committee.  
41

42 Approval was given by the Research and Governance leads for the three PCT areas  
43 where recruitment took place: Stockton-on-Tees; Redcar & Middlesbrough ; County  
44 Durham and Darlington.  
45  
46  
47

48 The RCGP Scientific Foundation Board awarded £ 3,850 to cover Transcription  
49 costs. No other funding was provided for the study.  
50  
51

52 There are no competing interests for any of the authors.  
53  
54

55 The authors agree to BMJOpen having exclusive licence to this original research.  
56  
57  
58  
59  
60

Table 1

Participant Number	Gender	Age	Salaried or Partner	Practice descriptor	Additional professional experience
01	F	50-59	S	Semi-rural Deprived	GP Postgraduate education
02	Male	50-59	S	Urban Deprived	Addiction medicine in primary care
03	Female	50-59	P	Urban Deprived; wealthy student population	Former Assoc. Specialist in CAMHS
04	Female	40-49	S	Semi-rural Deprived	Mental health Lead for a PCT
05	Female	20-29	S	Urban Deprived	
06	Male	40-49	P	Semi-rural Largely affluent	
07	Male	40-49	P	Semi-rural Mixed :	Child Protection Lead for a PCT
8	Female	30-39	S	Semi-rural Mixed :	
9	Male	50-59	P	Semi-rural Mixed :	GP lead for 'teen drop-in' clinic
10	Male	40-49	P	Urban Deprived	Mental Health and Child Protection Lead for a PCT. Substance misuse

11	Female	20-29	S	Urban Deprived	
12	Male	30-39	S	Semi-rural Mixed: largely affluent	
13	Female	30-39	S	Urban Deprived	
14	Male	40-49	P	Urban Deprived	
15	Male	40-49	P	Semi-rural Mixed :	
16	Female	20-29	S	Urban Deprived	
17	Male	30-39	S	Urban Deprived	
18	Female	40-49	P	Semi-rural Affluent	
19	Female	50-59	P	Semi-rural Mixed :	Child health lead



1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

Box 1 . Anxiety related to professional performance: In the consultation

*I'm quite anxious about mental health problems in young people cos I don't have a huge experience.....and I don't want to waste their (CAMHS practitioners') time (09;M;50-59;P)*

*I've always thought young people are challenging and still do and I have more questions than answers' (06;M; 40-49;P)*

*I think they are a difficult group...partly because of the way they present...and there should be lots of resources for them and there aren't so not knowing what to do is a bit of a theme really...the main anxiety is what to do.... (07;M; 40-49;P)*

*I find the adults will accept me at face value, generally. And they come with something usually fairly clear and they want that sorting out, it might not be straight forward, it might not even be simple they might have even brought things off the internet but it is a fairly clear baggage package... what I find with younger people with psychological or emotional disorders is it's not a clearly packed problem, it's in the extreme realms of the undefined. (06;M;40-49;P)*

Box 2. Anxiety related to professional performance: at an external level

*'...because it doesn't fit within any ticky box guidelines until time has passed I rarely know whether I've done the right thing, it's all in retrospect. (06;M;40- 49;P)*

*I'll bring people back in 1 week, I don't think this annoys my partners but it can become a bugbear....I'll squeeze them in when there are no appointments, which is probably making a rod for my own back and I wouldn't encourage trainees to do it, but I like the idea of seeing something through to its natural conclusion...its perhaps my own insecurity (14;M; 40-49;P)*

*What we don't have, in general practice is supervision...no counsellors are allowed to work without it but GPs are just sent out there, and I really do feel there is a huge need for it even if it is just one phone call-it's that ability to share the responsibility, not to dump it, but to genuinely share it. (04;F; 40-49;S)*



1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

Box 3. Anxiety related to professional performance: across disciplinary boundaries

*I know we don't meet all the people we refer to but we usually have something to hang it on because of our experience as junior doctors working in hospital but with CAMHS there is nothing. (08; F;30-39;S)*

*Some (mental health) creatures are on the verge of being mythical beasts, ...like psychotherapists,....educational psychologists (09,M;50-59;P)*

*CAMHS..it feels like it's a bit of a hotchpotch really, a patched together sort of service and I'm not sure who is control....people who counsel children-I don't know much about them, how much responsibility they take..... (07;M; 4049;M)*

Box 4. Anxiety related to interacting with young people

*Males with mental health issues worry me intensely, it really does seem that there is not an awful lot of trivia goes on there. By the time a male is presenting, because they don't have the tools to come to the GP very often, they don't understand that you can just come along when things are in their development, they usually come when something is really big ,black and bleak.(14;M;40-49;P)*

*I struggle a bit to work out how to word sort of mental health questions with to the sort of under 16 year olds particularly.... I suppose with adults I have my kind of, standard questions ..but using those sort of questions with young people often draws a blank face, and, so it's something I have to rephrase; I feel that I don't necessarily know their kind of lingo if you like...(17;M; 30-39;S)*

*Generally consulting with young people, I often find, if I'm being honest, probably more difficult than I would expect to find it. I think I probably have this unrealistic view of myself as really sort of approachable and you know still being quite young myself compared to other GPs, being able to communicate fairly easily and fairly well with young people, then always very quickly, it becomes apparent that actually no, you are a million miles away from where they are, and they don't really relate to you very well at all....(08; F; 30-39; S)*

Box 5. Anxiety associated with the complexity of presentations of adolescent emotional distress

*... you feel that there there are these big 'no go areas' in teenage consultations, around sex, drugs, alcohol... which loom over you like a black cloud and I'm thinking that they want to talk about it and I'm thinking that I want to talk about it but we can't talk about it... (11;F; 20-29; S)*

*They are missing school, in trouble with the police, youth offending team or not uncommonly the parent just comes by themselves...I never get a 14 year old acting out saying 'you know, I'm in trouble with my mam and dad' ( 10;M;40-49;P)*

*they are in absolute crisis one minute and then you see them a week later and they can hardly remember what it was all about (07;M; 40-49;P)*

*Its always a worry isn't it that you just completely get it wrong..I mean I'm conscious of this. I had someone in on Monday, parents, whose son had just hung himself at 21. I'd never seen him, he was a patient here. They had no idea anything was wrong. Nobody did....there is always that underlying things isn't there, that you might miss something catastrophic.. (01;F;50-59;S)*

#### Contributorship:

JR was the lead researcher and conducted all the interviews. She performed the primary analyses and is first author. AC and JF were involved with the study design from conception, met regularly throughout the analytical phase and commented on each draft of the manuscript.

## Reference list

1. Green H, McGinnity A, Meltzer H, Ford T, Goodman R. Mental Health of Children and Young People in Great Britain, 2004 : summary report. London: Office of National Statistics, 2005.
2. Singleton N, Bumpstead R, O'Brien M, Lee A, Meltzer H. Psychiatric morbidity in adults living in private households, 2000. Norwich: The Stationery Office, 2001.
3. Hawton K, Saunders KE, O'Connor RC. Self-harm and suicide in adolescents. *Lancet* 2012;379(9834):2373-82.
4. Kramer T, Garralda M. Child and adolescent mental health problems in primary care. *Advances in Psychiatric Treatment* 2000;6:287-94.
5. Hickie I, Fogarty A, Davenport T, Luscombe G, Burns J. Responding to experiences of young people with common mental health problems attending Australian general practice. *Medical Journal of Australia* 2007;187:S47-S52.
6. Gledhill J, Garralda M. The short-term outcome of depressive disorder in adolescents attending primary care: a cohort study. *Soc Psychiatry Psychiatr Epidemiol* 2011;46(10):993-1002.
7. Martinez R, Reynolds S, Howe A. Factors that influence the detection of psychological problems in adolescents attending general practices. *Br J Gen Pract* 2006;56(529):594-9.
8. Haller D, Sanci L, Sawyer S, Patton G. The identification of young people's emotional distress: a study in primary care. *Br J Gen Pract* 2009;59(560):e61-70.
9. Mauerhofer A, Berchtold A, Michaud PA, Suris JC. GPs' role in the detection of psychological problems of young people: a population-based study. *Br J Gen Pract* 2009;59(566):e308-14.
10. Churchill R. Child and Adolescent Mental Health. In: Cohen A, editor. *Delivering mental health for primary care: an evidence based approach*. London: RCGP, 2008:157-84.
11. Iliffe S, Gledhill J, da Cunha F, Kramer T, Garralda E. The recognition of adolescent depression in general practice: issues in the acquisition of new skills. *Primary Care Psychiatry* 2004;9(2):51-56.
12. Goldberg D, Huxley P. *Common mental disorders: a bio-social model*. London: Tavistock/Routledge, 1992.
13. Pilling S, Whittington C, Taylor C, Kendrick T. Identification and care pathways for common mental health disorders: summary of NICE guidance. *BMJ* 2011;342:d2868.
14. Dowrick C. *Beyond Depression: a new approach to understanding and management*. 2nd ed. Oxford: Oxford University Press, 2009.
15. Reeve J. Interpretive medicine: Supporting generalism in a changing primary care world. *Occas Pap R Coll Gen Pract* 2010(88):1-20, v.
16. Dowrick C, May C, Richardson M, Bundred P. The biopsychosocial model of general practice: rhetoric or reality? *Br J Gen Pract* 1996;46(403):105-7.
17. Karasz A, Dowrick C, Byng R, Buszewicz M, Ferri L, Hartman TCO, et al. What we talk about when we talk about depression: doctor-patient conversations and treatment decision outcomes. *Br J Gen Pract* 2012;62(594):e55-e63.
18. Kennedy I. Getting it right for children and young people: overcoming cultural barriers in the NHS so as to meet their needs. A review by Sir Ian Kennedy. London: Department of Health, 2010.
19. Layard R. How mental health loses out in the NHS. London: London School of Economics and Political Science, 2012.
20. Department for Education and Skills. Every Child Matters: change for children London: HMSO, 2002.
21. Department of Health. *Child and Adolescent Mental Health-National Service Framework for Children, Young People and Maternity Services* London: Department of Health 2004.
22. Department of Health. *Healthy Lives, Brighter Futures* London: The Stationery Office, 2009.

1. Green H, McGinnity A, Meltzer H, Ford T, Goodman R. Mental Health of Children and Young People in Great Britain, 2004 : summary report. London: Office of National Statistics, 2005.
2. Singleton N, Bumpstead R, O'Brien M, Lee A, Meltzer H. Psychiatric morbidity in adults living in private households, 2000. Norwich: The Stationery Office, 2001.
3. Hawton K, Saunders KE, O'Connor RC. Self-harm and suicide in adolescents. *Lancet* 2012;379(9834):2373-82.
4. Kramer T, Garralda M. Child and adolescent mental health problems in primary care. *Advances in Psychiatric Treatment* 2000;6:287-94.
5. Hickie I, Fogarty A, Davenport T, Luscombe G, Burns J. Responding to experiences of young people with common mental health problems attending Australian general practice. *Medical Journal of Australia* 2007;187:S47-S52.
6. Gledhill J, Garralda M. The short-term outcome of depressive disorder in adolescents attending primary care: a cohort study. *Soc Psychiatry Psychiatr Epidemiol* 2011;46(10):993-1002.
7. Martinez R, Reynolds S, Howe A. Factors that influence the detection of psychological problems in adolescents attending general practices. *Br J Gen Pract* 2006;56(529):594-9.
8. Haller D, Sanci L, Sawyer S, Patton G. The identification of young people's emotional distress: a study in primary care. *Br J Gen Pract* 2009;59(560):e61-70.
9. Mauerhofer A, Berchtold A, Michaud PA, Suris JC. GPs' role in the detection of psychological problems of young people: a population-based study. *Br J Gen Pract* 2009;59(566):e308-14.
10. Churchill R. Child and Adolescent Mental Health. In: Cohen A, editor. *Delivering mental health for primary care: an evidence based approach*. London: RCGP, 2008:157-84.
11. Iliffe S, Gledhill J, da Cunha F, Kramer T, Garralda E. The recognition of adolescent depression in general practice: issues in the acquisition of new skills. *Primary Care Psychiatry* 2004;9(2):51-56.
12. Goldberg D, Huxley P. *Common mental disorders: a bio-social model*. London: Tavistock/Routledge, 1992.
13. Dowrick C. *Beyond Depression: a new approach to understanding and management*. 2nd ed. Oxford: Oxford University Press, 2009.
14. Reeve J. Interpretive medicine: Supporting generalism in a changing primary care world. *Occas Pap R Coll Gen Pract* 2010(88):1-20, v.
15. Dowrick C. Reasons to be cheerful ? Reflections on GPs' responses to depression. *Br J Gen Pract* 2009;59(566):636-7.
16. Dowrick C, May C, Richardson M, Bundred P. The biopsychosocial model of general practice: rhetoric or reality? *Br J Gen Pract* 1996;46(403):105-7.
17. Karasz A, Dowrick C, Byng R, Buszewicz M, Ferri L, Hartman TCO, et al. What we talk about when we talk about depression: doctor-patient conversations and treatment decision outcomes. *Br J Gen Pract* 2012;62(594):e55-e63.
18. Kennedy I. Getting it right for children and young people: overcoming cultural barriers in the NHS so as to meet their needs. A review by Sir Ian Kennedy. London: Department of Health, 2010.
19. Layard R. How mental health loses out in the NHS. London: London School of Economics and Political Science, 2012.
20. Department for Education and Skills. Every Child Matters: change for children London: HMSO, 2002.
21. Department of Health. *Child and Adolescent Mental Health-National Service Framework for Children, Young People and Maternity Services* London: Department of Health 2004.
22. Department of Health. *Healthy Lives, Brighter Futures* London: The Stationery Office, 2009.
23. National Institute for Health & Clinical Excellence. Depression in children and young people: identification and management in primary, community and secondary care. *National Clinical Practice Guidelines*. London: NICE, 2005.

- 1  
2  
3 24. Tait L. To disclose or not to disclose psychological problems to GPs. *Br J Gen Pract*  
4 2009;59(566):638-9.  
5 25. Biddle L, Donovan JL, Gunnell D, Sharp D. Young adults' perceptions of GPs as a help source for  
6 mental distress: a qualitative study. *Br J Gen Pract* 2006;56(533):924-31.  
7 26. Glaser B, Strauss A. *The discovery of grounded theory*. Chicago: Aldine, 1967.  
8 27. Charmaz K. *Constructing Grounded Theory: A practical guide through qualitative analysis*.  
9 Thousand Oaks, California: Sage Publications, Inc, 2006.  
10 28. Clarke A. *Situational analysis: Grounded Theory after the postmodern turn*. Sage Publications, Inc.  
11 Thousand Oaks, California: Sage Publications, Inc, 2005.  
12 29. Chew-Graham CA, May CR, Perry MS. Qualitative research and the problem of judgement:  
13 lessons from interviewing fellow professionals. *Family Practice* 2002;19(3):285-89.  
14 30. Heath I. Uncertain clarity: contradiction, meaning, and hope. *Brit J Gen Pract* 1999;49(445):651-  
15 57.  
16 31. Schon D. *The reflective practitioner*. San Francisco: Jossey-Bass, 1983.  
17 32. Iliffe S, Williams G, Fernandez V, Vila M, Kramer T, Gledhill J, et al. Treading a fine line: is  
18 diagnosing depression in young people just medicalising moodiness? *Br J Gen Pract*  
19 2009;59(560):156-7.  
20 33. Iliffe S, Gallant C, Kramer T, Gledhill J, Bye A, Fernandez V, et al. Therapeutic identification of  
21 depression in young people: lessons from the introduction of a new technique in general  
22 practice. *Br J Gen Pract* 2012;62(596):e174-82.  
23 34. Sawyer SM, Afifi RA, Bearinger LH, Blakemore SJ, Dick B, Ezech AC, et al. Adolescence: a  
24 foundation for future health. *Lancet* 2012;379(9826):1630-40.  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

## Appendix 1 Early Topic Guide

1. I'd like to talk about your experiences of consulting with young people in general
  - How do you find this age group?
  - Is it very different to consulting with older patients?
  - What sort of problems do you see? Do they consult often?
2. Can we talk more about consulting with young people who may have psychological/mental health problems
  - How do you find this clinical area?
  - What about seeing YP alone/ with 'another'?
  - Any areas particularly tricky to broach ?
3. How do you consider possible 'mental health problems' which presenting in young people ?
  - Do any examples come to mind ?
  - What approach did you take
  - What worked well? What was difficult?
  - Is it different with other age groups
4. What are your thoughts on 'depression' and 'anxiety' in young people ?
  - Do you see much of it?
  - Does this differ from other age groups?
  - What options are there in primary care?
5. Do you think GPs have a role/or not in promoting emotional well-being in young people? Explore

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

Research checklist

As this is a qualitative study it does not fall within the parameters of the recommended research checklists.

A statement to this effect is included in the covering letter.

For peer review only



1  
2  
3  
4  
5  
6  
7  
8  
9 **“.. I think this is maybe our Achilles Heel...” Exploring GPs’ responses to**  
10 **young people presenting with emotional distress in general practice. A**  
11 **qualitative study.**  
12

13  
14  
15  
16 Jane H Roberts , Ann Crosland, John Fulton

17  
18 Department of Pharmacy, Health & Well-being, Faculty of Applied Sciences, The  
19 Science Complex, Wharnclyffe St, Sunderland, SR1 3SD.

20  
21 Jane H Roberts, Clinical Senior Lecturer, University of Sunderland,

22  
23 Ann Crosland, Professor of Nursing , University of Sunderland,

24  
25 John Fulton, Principal Lecturer, University of Sunderland,  
26  
27

28  
29 Correspondence to Jane H Roberts

30  
31 [jane.roberts@sunderland.ac.uk](mailto:jane.roberts@sunderland.ac.uk)  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60



1  
2  
3  
4  
5  
6  
7 “.. I think this is maybe our Achilles Heel....” Exploring GPs’ responses to  
8 young people presenting with emotional distress in general practice. A  
9 qualitative study.  
10

## 11 Article summary

### 12 Article Focus:

- 13 1. an exploratory study
- 14 2. to examine GPs’ views and experiences of consulting with young people  
15 experiencing emotional distress
- 16 3. to better understand GPs perspectives

### 17 Key Messages

- 18 1. ~~GPs collectively describe a~~ Anxiety about practice experienced when  
19 consulting with young people and uncertainty ~~is the dominant finding in a first~~  
20 stage analysis of a qualitative study. about their clinical practice when  
21 consulting with young people in distress, This is independently of age and  
22 gender of GP
- 23 2. Anxiety relates to professional performance; interacting with young  
24 people and the complex nature of presentations of emotional distress in  
25 primary care
- 26 3. Unless anxiety and related uncertainty ties about practice are  
27 addressed GPs will continue to miss opportunities to address early  
28 emotional difficulties and young people’s mental health needs in primary care  
29 will continue to be poorly met

### 30 Strengths and Limitations

- 31 1. Qualitative research in under -examined areas offers new insights  
32 and explores why behaviours might arise
- 33 2. The data presented contributes to ~~to~~ theory building and offers  
34 theoretical generalizability
- 35 3. Theoretical sampling led to only white British born GPs participating  
36 so other cultural perspectives were not included

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11 **“.. I think this is maybe our Achilles Heel....” Exploring GPs’ responses to**  
12 **young people presenting with emotional distress in general practice. A**  
13 **qualitative study.**  
14

15  
16  
17  
18 **Main text**

19  
20 **Introduction**

21  
22 Emotional distress in young people is common. It may be the affective response to  
23 the challenges of everyday life or may indicate a mental health disorder compatible  
24 with a psychiatric diagnosis ~~an associated mental health problem~~. The most recent  
25 and widely cited household survey reports ~~, with~~ at least 10% of 10-15 year olds  
26 affected,<sup>1</sup> and 17 % of 16-19 year olds <sup>2</sup> ~~(based on household surveys to )~~ to have  
27 symptoms consistent with a mental health disorder as defined by the ICD-10.

28  
29 Behavioural manifestations of emotional distress might include ~~Proxy markers of~~  
30 distress, such as reported incidences of self-harm which, at a conservative estimate,  
31 appears to affect around 10% of adolescents, as reported in six studies cited by  
32 Hawton et al in a recently published review ~~derived from community based studies,~~  
33 show 10% of adolescents report having self-harmed.<sup>3</sup>

34  
35 Data from populations of young people who consult their GP reveal higher rates of  
36 psychological distress, of the order of 20-30%.<sup>4,5</sup> GPs identify serious mental illness  
37 but often fail to detect less severe manifestations<sup>6</sup> and appear reluctant to discuss  
38 emotional issues<sup>7</sup>; unless offered cues by the young person in the consultation<sup>8</sup> or if  
39 other factors are present such as a previous history of a suicide attempt or a pattern  
40 of frequent consulting<sup>9</sup>. Young people’s presentations in primary care are often  
41 complex and present with behavioural, psychosocial, academic and familial  
42 problems which can be problematic to untangle in contrast to adult mental health  
43 manifestations which, although variable, may be less intense in their presentation.  
44 They Adolescent emotional distress may suggest indicate underlying co-morbid  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

Field Code Changed

Field Code Changed

Field Code Changed

Field Code Changed

Field Code Changed

Field Code Changed

Field Code Changed

Field Code Changed

1  
2  
3  
4  
5  
6  
7 mental health problems. ~~and it~~ It has been ~~reported suggested~~ that often the 'most  
8 important features in terms of assessment may be concealed or hidden',<sup>10</sup>.

Field Code Changed

9  
10 A key concern is the difficulty of distinguishing between 'moodiness' or a persisting  
11 emotional disorder and GPs have expressed a worry at 'over-medicalising young  
12 people's lives'.<sup>11</sup> Illiffe & colleagues found that GPs were uncomfortable about  
13 making a diagnosis of depression in young people (the most common, but often  
14 coexisting, mental health problem in adolescence ).

Field Code Changed

15  
16  
17  
18 ~~This sits in contrast to~~ On the other hand GPs' are increasingly involved ment of in  
19 managing common mental health problems in older patients.<sup>12 13</sup> ~~and also to a~~  
20 ~~broadening of the frames of reference by which emotional distress in adults is~~  
21 ~~regarded~~. Although a biomedical perspective dominates, supported by an array of  
22 NICE clinical guidelines, alternative frameworks for considering adult mental health  
23 problems have been offered.<sup>13 14</sup> Dowrick and Reeve<sup>14 15</sup> ~~have offered alternative~~  
24 ~~frameworks and~~ refer to the insights derived from the wisdom traditions in informing  
25 their work which moves away from a positivist understanding of emotional distress to  
26 an approach which incorporates ideas of personal agency and encourages hope.<sup>15</sup>

Field Code Changed

Field Code Changed

Field Code Changed

Formatted: Superscript

27  
28 Historically, research has found GPs ~~have been found~~ to be largely dismissive of  
29 their role in addressing social issues in adult mental ill-health.<sup>16</sup> Contemporary  
30 studies reveal a shift although this position is shifting with greater awareness of the  
31 lay perspective, which typically favours the a social model causes of adult mental  
32 ill-health ~~(notably depression) as being social in origin~~.<sup>17</sup> and a matched response  
33 by GPs mirroring popular social constructions of distress.

Field Code Changed

Field Code Changed

34  
35  
36  
37  
38  
39  
40  
41  
42 Despite the challenge of responding to emotional distress in adolescence and the  
43 patchy, often inadequate provision of secondary care services,<sup>18 19</sup> a series of policy  
44 directives have emphasised the role of GPs and other front-line services, in the  
45 promotion of psychological well-being and the early indication of difficulties.<sup>20 21 22</sup>

Field Code Changed

Field Code Changed

Field Code Changed

46 Practitioners are expected to have 'sufficient knowledge, training and support 'in this  
47 area including competence in 'active listening' and conversational technique',<sup>23</sup>.

Field Code Changed

48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
There is a growing body of evidence examining young people's experiences of  
talking to GPs about emotional problems. They reveal a mixed picture including a  
reluctance to disclose<sup>24</sup>, a fear of being judged or offered medication<sup>25</sup>. Much less

Field Code Changed

Field Code Changed

is known about GP perspectives. This paper presents a qualitative, exploratory study which examines GPs' views and experiences of consulting with young people presenting with emotional distress.

## Method

### Study Design

The study took place in the North East of England in 18 general practices based in urban, rural and semi-rural communities serving predominantly socio-economically disadvantaged patients. The qualitative study comprised of in depth individual interviews with GPs recruited using theoretical sampling. As early theoretical ideas emerged successive GPs were recruited on the basis of their capacity to contribute to the development or abandonment of initial theoretical constructs.

Data were collected between January 2010 to May 2011

### Participants

GPs with less than four years clinical experience were excluded. The initial recruits were selected on the basis of their relevant experience and their ability to generate early data which would scope the terrain of the area under enquiry; for example having a role as mental health lead or previous experience working in Child & Adolescent Mental Health services (CAMHS)

GPs were approached by telephone and email contact and sent information sheets. A follow-up contact established their verbal consent to meet at a location of their choice. Two GPs approached declined to participate. One cited forthcoming extended annual leave and another a view that as the senior partner he saw relatively few younger aged patients and suggested recruitment of a younger GP in the same practice.

Ethical approval by the Local Research Ethics Committee, the seven Primary Care Trust organizations of the region and the University of Sunderland was granted before data collection began.

### Data collection and analysis

1  
2  
3  
4  
5  
6  
7 The audio-taped semi-structured interviews were transcribed verbatim with consent.  
8 An initial topic guide was used with the first tranche of participants based on the  
9 extant literature and developed through discussion. The topic guide was then revised  
10 on the basis of ideas arising from the early interviews, and the iterative analysis  
11 which began as soon as the first interview was undertaken. The interview guides  
12 explored doctors' experiences of consulting with young people in general and those  
13 presenting with psychological or mental health problems, GPs' understanding of  
14 depression and anxiety in adolescence, of how emotional distress presents in the  
15 surgery and the role of the GP in promoting emotional well-being in young people  
16 (See appendix 1). The guide was refined to include questions about how structural  
17 changes impacted on, and consultation style shaped, practice.

18  
19  
20  
21  
22  
23 The interviews lasted between 50 to 75 minutes. Field notes and theoretical memos  
24 were kept throughout the period of data collection and analysis.

25  
26  
27 The transcripts were coded and analysed using the grounded theory method  
28 described by Strauss and Glaser<sup>26</sup> and revised by Charmaz<sup>27</sup>. The constant  
29 comparative method of analysis is core to the process and informs the theoretical  
30 sampling of recruits. Early ideas were tested with subsequent participants and found  
31 to be either substantiated or rejected through the iterative process of constant  
32 comparison supported by theoretical sampling. Situational maps, both 'messy' and  
33 'ordered', were constructed during this phase of analysis.<sup>28</sup>

34  
35  
36  
37  
38 The data presented here ~~wasere produced generated~~ after the first level of analysis  
39 was completed, during which only the open codes were iteratively developed by JR  
40 and subjected ed to further examination by AC (primary care academic) and  
41 JF(sociologist). Further analysis of the axial and selective codes will be presented in  
42 two subsequent companion papers.

## 43 44 45 Results

46  
47  
48 Nineteen GPs participated, 10 women. (Table1). The early iterative analysis of the  
49 data found the open codes to support a dominant narrative of anxiety ~~and~~  
50 ~~uncertainty about practice~~ under-pinning the majority of the research interviews. This  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

Field Code Changed

Field Code Changed

Field Code Changed

pervasive and disabling emotional response to encounters with emotionally distressed young people appeared to coalesce around three domains.

These can be viewed as anxiety ~~and uncertainty~~ experienced by GPs in response to:

- 1) professional performance; in the consultation, at an external level, across disciplinary boundaries;
- 2) interacting with young people; and
- 3) the complexity of presentations of adolescent emotional distress

Each of the three themes will be presented in turn and supported by illustrative quotations taken from the transcripts (see boxes 1-6). GP participants are identified by identifier number, gender, age range and whether salaried or a partner ( as presented in Table 1.)

**1. i. Anxiety related to professional performance: operating *In the consultation***

A coherent narrative emerged, gathered from almost all of the participants, of practitioners being anxious in the consultation because of an uncertainty about what to do and of what was expected of them as primary care clinicians.

~~A prevailing finding was the~~ This resulted in a sense of professional impotence ~~which was associated with seeing or suspecting emotional distress in this age group.~~ It was acknowledged that feeling uncertain about how best to proceed, and unsure of practice, led to a sense of disempowerment -through not knowing what to do;. This was -in contrast to accounts of working with older patients where the options for GPs appear more clearly defined. The data generated collected by the open code analysis suggested that not being able to formulate the initial presentation by a young person into a definable 'disorder' created a sense of operating in uncharted territory.

~~This Anxiety~~ was amplified by the lack of exposure to adolescent mental health in undergraduate medical education which was the ~~unanimously experience shared by~~ of all participants. Where the topic had been included in the curriculum, it was often

7

**Formatted:** Numbered + Level: 1 +  
Numbering Style: 1, 2, 3, ... + Start at: 1 +  
Alignment: Left + Aligned at: 0.25" + Indent  
at: 0.5"

**Formatted:** Font: Not Bold

**Formatted:** Indent: Left: 0.25"

**Formatted:** Font: Not Bold

restricted to severe mental disorder for example being assigned to medical teams looking after such as adolescents hospitalized with anorexia nervosa.

(See box 1).

**1. ii Anxiety related to professional performance: operating at an external level**

The ~~A~~ lack of benchmarks in practice meant assessing one's performance in relation to peers was problematic since no 'gold standard' existed. The only NICE guideline which was referenced (concerning the management of depression in under 18 year olds) was regarded as having "hampered GPs" from becoming involved in the management of adolescent depression since the Guideline did not advocate the use of anti-depressants and, with access to psychological therapies piecemeal, appeared to and supporting a position view that there was little to be offered in primary care.

Constraints in practice led to frustration and an anxiety about management. For example, Varying arrangements within practices governing access to appointments and the ease, or not, of maintaining continuity of care were seen to contribute to professional anxiety by impeding attentive 'watchful waiting' and some GPs described attempts to circumvent inflexible appointment systems in order to be more available to patients.

A lack of professional supervision was identified by a small number of more experienced GPs involved with Postgraduate Training and provision of mental health services at a regional level, and contrasted to systems for other professionals working with emotionally distressed patients. Leaving GPs to rely on their own personal resources, on informal collegiate support or ad hoc relationships with colleagues in secondary care resulted in a fragile structure which could amplify rather than ameliorate anxiety.

(See box 2).

**1. iii Anxiety related to professional performance: across disciplinary boundaries**

Formatted: Numbered + Level: 1 +  
Numbering Style: 1, 2, 3, ... + Start at: 1 +  
Alignment: Left + Aligned at: 0.25" + Indent  
at: 0.5"

Formatted: Font: Not Bold



1  
2  
3  
4  
5  
6  
7 GPs across the board expressed frustration with access to secondary care services,  
8 reporting long delays and frequent rejection of referrals, and a lack of clarity about  
9 how the services were structured and governed. GP experiences and degrees of  
10 frustration varied with an emerging picture of problematic access to services being  
11 associated with higher levels of professional anxiety. Where GPs described Mmore  
12 constructive cross-disciplinary relationships were described with CAMHS workers  
13 practitioners offering clinical updates meetings, and were consultants were  
14 accessible by telephone, less anxiety was voiced.

15  
16  
17  
18  
19 Unfamiliarity with the roles and responsibilities of CAMHS practitioners, coupled with  
20 an obligation to refer in the absence of other options, left some GPs feeling uncertain  
21 about the clinical care pathway and unsure about practice.

22  
23  
24 (See box 3.)

## 26 **2. Anxiety related to interacting with young people**

27  
28 The open codes showed a dominant finding of GPs expressing The early finding of  
29 anxiety and uncertainty in this area was under-pinned by associated with the  
30 difficulties GPs talked about experiencing when communicating with young people  
31 in general. Neither the age nor the gender of the GP appeared to facilitate  
32 communication, with younger and female GPs similarly as uneasy as older male and  
33 female GPs. Female patients were generally considered to be easier to talk with  
34 whilst young men were seen to be more challenging because of their perceived  
35 reluctance to seek help and their tendency to present late.

36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
Communication difficulties included establishing a rapport, finding the right words  
and tone to use and dealing with silence. An inability to read the non-verbal signs,  
and to translate an often terse description from the young person into a coherent  
picture of their internal emotional state, left many GPs either relying on the  
accompanying parent or closing down the consultation. Being able to find common  
ground was identified as being key to beginning the process of establishing rapport.

Young people were seen as a highly heterogeneous group who showed variability  
from one presentation to the next (intra-variability), and also across lines of age and  
gender (inter-variability). Knowing what was 'normal' for an individual, particularly if it

9

Formatted: Numbered + Level: 1 +  
Numbering Style: 1, 2, 3, ... + Start at: 1 +  
Alignment: Left + Aligned at: 0.25" + Indent  
at: 0.5"



was presented as a principal reason for consulting with the GP, was perceived as problematic and anxiety provoking, both for the young person and for the GP.

(See box 4)

### 3. Anxiety associated with the complexity of presentations of adolescent emotional distress

GPs' accounts of their experiences consulting with young people experiencing distress described a terrain beset with pitfalls, associated with the unspoken or with complex narratives embedded in social contexts; and presented in an undifferentiated form. GPs spoke of There was a sense of unpredictability and volatility to presentations which left GPs them -uncertain about how the patient narrative might unfold and how much input to offer at the initial consultation. This was in contrast. In particular this generated anxiety associated with ~~to~~ the rare but grave consequences which might arise when a young person seriously attempted or completed suicide; a clinical experience to which many GPs referred and which could lead to enduring professional anxiety. (See box 5)

Although it was accepted that uncertainty as a feature of general practice was not restricted to the clinical area of youth mental health, the early first stage analysis showed a distinct narrative emerging in which adolescent mental health was seen as more notably anxiety provoking because of its more nebulous presentation and multiple confounding factors, which largely pertained ing to the social environment. The account given in the consulting room was described as the 'iceberg' indicating that often much is left hidden, or unsaid, but which nevertheless has to be raised at some point if the young person's distress is to be addressed.

Not only is there a dominant narrative of anxiety and uncertainty surrounding how GPs make sense of adolescent emotional distress, but similar responses surround are associated with management options. Few GPs expressed any degree of confidence about how they would tackle individual presentations. A small number of those with additional roles in mental health or working with patients with substance abuse problems spoke of a more systematic approach to organizing and offering care. However but even established GPs with personal experience of working in 'a

Formatted: Numbered + Level: 1 +  
Numbering Style: 1, 2, 3, ... + Start at: 1 +  
Alignment: Left + Aligned at: 0.25" + Indent  
at: 0.5"

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

teen drop-in clinic' or with drug dependent patients ~~were described uncertainty about uncertain of~~ their practice. A paucity of treatment options was a consistent re finding along with a lack of clarity about what GPs might reasonably be expected to do, if supported by adequate professional development.

## Discussion

### Summary

Anxiety ~~and uncertainty about practice, coupled with a perceived reduced range of options and lack of clarity of expectations,~~ associated with diverse presentations of adolescent emotional distress in primary care, emerged from all GP participant accounts in the first stage of analysis, and from the early iterative analysis of the data. Anxiety was associated with the clinical consultation, with what was expected of the GP, and how they might best respond in the absence of few clinical guidelines and limited options to involve other health and social care professionals. Unease when communicating with young people and ~~of difficulties~~ interpreting their accounts of distress inhibited GPs, and This was compounded by the complexity of presentations which ranged from familial discord to school refusal to offending behaviour, usually in the absence of any clear diagnosis. The heterogeneity of adolescent behaviour taxed GPs as did the unpredictability of the unfolding clinical presentation which ~~might could~~ settle spontaneously or might develop into a serious mental health disorder.

Whilst there was a spectrum of levels of anxiety experienced by GPs, there was a prevailing universality about the experience. How GPs responded and managed the perceived threat to professional competence and confidence was interrogated in the next stage of the analysis which would lead to the development of the axial codes, or pillars, of the emerging conceptual model (presented elsewhere).

### *Strengths and limitations*

The management of adolescent mental health problems remains an under-investigated area of clinical practice. Previous research studies have largely often been conducted by psychiatrists whose perspective is different to that of GPs

~~responding to undifferentiated distress in the consulting room and whilst plurality of perspectives is important, unless more is known and understood about how GPs perceive the area many assumptions will go unchallenged.~~ Using grounded theory, augmented by situational analysis, permits a rich exploration of the territory and facilities theory building.

Theoretical sampling supports theory development whilst not purporting to provide universal generalizability. After 19 in-depth interviews, buttressed by situational analysis, no new themes emerged and theoretical saturation was reached. All of the respondents were white British and whilst they were recruited on the basis of their contribution to the study, it must be acknowledged that the absence of including the experiences of GPs raised and educated in different cultural contexts will lead to the silencing of other cultural perspectives.

The lead researcher and interviewer is a GP (JR). Interviewing peers has been described<sup>29</sup> as enriching the data collection because of the shared knowledge and familiarity with the clinical territory but it can lead to collusion between interviewer and respondent which needs attention and reflexivity. Co-contributors AC and JF have academic expertise in social policy and sociology which strengthened the analysis.

#### *Comparison with existing literature*

Heath asserts that *a commitment to uncertainty is fundamental to general practice*:<sup>30</sup>: 651) and Schon has described this operative landscape as a 'swampy lowland' proposing a model which advocates reflective practice as the key to dealing with uncertainty.<sup>31</sup> A quest for certainty in areas of complex practice, especially when it concerns individual experiences can be counter-productive and scholars have cautioned against clinging to the 'shelter of diagnosis'<sup>1532</sup> when what is required involves attention to alleviating suffering and working purposefully with patients to catalyse their own creative capacity.<sup>1415</sup> Illife et al's earlier cited work demonstrated that when GPs were fixed on the concept of depression as disease they were uncomfortable talking to young people.<sup>3233 3334</sup>

Field Code Changed

Field Code Changed

Field Code Changed

Field Code Changed

Field Code Changed

Field Code Changed

Field Code Changed

1  
2  
3  
4  
5  
6  
7 This study which contributes to building a theoretical model, suggests that ~~it is the~~ anxiety and perceived -threats to professional competence, can be experienced at  
8 multiple levels, and are amplified with regard to the complexity of adolescent  
9 presentations. ~~This and perceived paucity of management options which can~~  
10 ~~compromise GP's~~ professional engagement ~~and inhibits them from taking a~~  
11 ~~more active role with young people. Understanding more about why some GPs can~~  
12 creatively respond to the anxiety and lack of certainty about expectations defines  
13 the next stage of the analysis.  
14  
15  
16  
17  
18

#### 19 *Implications for practice and research*

20  
21 Inadequate educational preparation, both at under and post-graduate level, is pivotal  
22 in failing to address sustaining the anxiety around clinical practice. Doctors need to  
23 be introduced to the developmental trajectory of adolescence and the conceptual  
24 framework which locates adolescence as the foundation of future health <sup>3436</sup> both in  
25 undergraduate education and revisited in continuing professional development .  
26  
27

Field Code Changed

28 This approach will help GPs to understand more about why addressing emotional  
29 distress in the second decade of life is important.  
30  
31

32 GPs need good quality educational exposure and preparation to deal with the multi-  
33 axial development of adolescence and the emergence of mental health disorders in  
34 the 10 to 20 year olds. The current psychiatric classification systems do not facilitate  
35 clinical practice in this domain at primary care level.  
36  
37

38  
39 In addition, the links between general practice and CAMHS need to be strengthened  
40 both in terms of education and understanding more of how each discipline operates,  
41 but also at a pragmatic, operational level. If cross-disciplinary practice was  
42 facilitated more treatment options would be presented at a primary care or early  
43 intervention level .  
44

45  
46 More research is needed to demonstrate Eevidence of effective, feasible, primary  
47 care based brief behavioural interventions which would equip GPs to engage with  
48 young people with greater confidence and support the development of evidence  
49 based policy.  
50  
51  
52  
53  
54

1  
2  
3  
4  
5  
6  
7 At a systemic level, this study shows that external factors are important in influencing  
8 practice and can moderate or exacerbate levels of anxiety. Systems which improve  
9 access to care for young people need to be introduced at practice level and  
10 supported by policy.

11  
12  
13  
14  
15 ~~The needs of young people are ill-served by the current provision<sup>18,19</sup> and whilst~~  
16 ~~rhetoric has called for GPs to be more involved, unless we address the disabling~~  
17 ~~anxiety and uncertainty in this area practice will remain static with GPs reluctant to~~  
18 ~~become involved in youth mental health.~~

#### How this fits in

19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

GPs are known to have difficulty recognizing and responding to adolescent emotional distress. Reluctance to medicalize distress has been reported.

This study shows that anxiety ~~and uncertainty~~ about practice in this complex clinical area ~~are~~ universal and independent of age, gender, ~~and~~ level of experience of GP.

If GPs are to play a more active role in the early identification and intervention of distress we need to know more about ~~the factors which ameliorate or exacerbate professional anxiety about practice, what promotes or inhibits professional anxiety and facilities~~ greater GP engagement with young people

Critically, adolescent mental health needs to feature in undergraduate and postgraduate curricula.

Table 1

Participant Number	Gender	Age	Salaried or Partner	Practice descriptor	Additional professional experience
01	F	50-59	S	Semi-rural	GP Postgraduate education
02	Male	50-59	S	Deprived Urban Deprived	Addiction medicine in primary care
03	Female	50-59	P	Urban Deprived; wealthy student population	Former Assoc. Specialist in CAMHS
04	Female	40-49	S	Semi-rural Deprived	Mental health Lead for a PCT
05	Female	20-29	S	Urban Deprived	
06	Male	40-49	P	Semi-rural Largely affluent	
07	Male	40-49	P	Semi-rural Mixed :	Child Protection Lead for a

15

					PCT
8	Female	30-39	S	Semi-rural	
9	Male	50-59	P	Mixed : Semi-rural	GP lead for 'teen drop-in' clinic
10	Male	40-49	P	Mixed : Urban  Deprived	Mental Health and Child Protection Lead for a PCT. Substance misuse
11	Female	20-29	S	Urban	
12	Male	30-39	S	Deprived Semi-rural	
13	Female	30-39	S	Mixed: largely affluent Urban	
14	Male	40-49	P	Deprived Urban	
15	Male	40-49	P	Deprived Semi-rural	
16	Female	20-29	S	Mixed : Urban	
17	Male	30-39	S	Deprived Urban	
18	Female	40-49	P	Deprived Semi-rural	
19	Female	50-59	P	Affluent Semi-rural Mixed :	Child health lead

## Anxiety paper Boxes i

Box 1. Anxiety related to professional performance: In the consultation

*I've always thought young people are challenging and still do and I have more questions than answers' (06;M; 40-49;P)*

*so not knowing what to do is a bit of a theme really (07;M; 40-49;P)*

*I find the adults will accept me at face value, generally. And they come with something usually fairly clear and they want that sorting out, it might not be straight forward, it might not even be simple they might have even brought things off the internet but it is a fairly clear baggage package... what I find with younger people with psychological or emotional disorders is it's not a clearly packed problem, it's in the extreme realms of the undefined. (06;M;40-49;P)*

*I know we don't meet all the people we refer to but we usually have something to hang it on because of our experience as junior doctors working in hospital but with CAMHS there is nothing. (08; F;30-39;S)*

Box 2 Anxiety related to professional performance: at a structural level

*NICE guidelines a few years ago looked at depression in young people and kind of hampered our ability to do anything with them really (07;M; 40-49;P)*

*'...because it doesn't fit within any ticky box guidelines until time has passed I rarely know whether I've done the right thing, it's all in retrospect. (06;M;40- 49;P)*

*I'll bring people back in 1 week, I don't think this annoys my partners but it can become a bugbear....I'll squeeze them in when there are no appointments, which is probably making a rod for my own back and I wouldn't encourage trainees to do it, but I like the idea of seeing something through to its natural conclusion...its perhaps my own insecurity (14;M; 40-49,P)*

*What we don't have, in general practice is supervision...no counsellors are allowed to work without it but GPs are just sent out there, andI really do feel there is a huge need for it even if it is just one phone call-it's that ability to share the responsibility, not to dump it, but to genuinely share it.(04;F; 40-49;S)*

**Comment [J1]:** i need to change structural to extenal



## Anxiety Boxes ii

## Box 3. Anxiety related to professional performance: across disciplinary boundaries

*Some (mental health) creatures are on the verge of being mythical beasts, ...like psychotherapists,....educational psychologists (09,M;50-59;P)*

*CAMHs..it feels like it's a bit of a hotchpotch really, a patched together sort of service and I'm not sure who is control....people who counsel children-I don't know much about them, how much responsibility they take.. (07;M; 4049;M)*

## Box 4. Anxiety related to interacting with young people

*Generally consulting with young people, I often find, if I'm being honest, probably more difficult than I would expect to find it. I think I probably have this unrealistic view of myself as really sort of approachable and you know still being quite young myself compared to other GPs, being able to communicate fairly easily and fairly well with young people, then always very quickly, it becomes apparent that actually no, you are a million miles away from where they are, and they don't really relate to you very well at all....(08; F; 30-39; S)*

*Males with mental health issues worry me intensely, it really does seem that there is not an awful lot of trivia goes on there. By the time a male is presenting, because they don't have the tools to come to the GP very often, they don't understand that you can just come along when things are in their development, they usually come when something is really big ,black and bleak.(14;M;40-49;P)*

*I struggle a bit to work out how to word sort of mental health questions with to the sort of under 16 year olds particularly.... I suppose with adults I have my kind of, standard questions ..but using those sort of questions with young people often draws a blank face, and, so it's something I have to rephrase; I feel that I don't necessarily know their kind of lingo if you like...(17;M; 30-39;S)*

*So he went off to do a urine sample and I was pleased to speak to his parents without him, seemed easier to talk about some of the mental issues without him there... (017;M; 30-39;S)*

*With children and teenagers it tends to be you controlling the pace of the consultation.... and you finish the consultation when you want to (07;M; 40-49;P)*

## Anxiety Box iii

## Box 4. Anxiety related to interacting with young people

*there's no such thing as a typical teenage presentation ( 13;F; 30-29;S)*

*...and her mum said, as mum's do, oh she has dreadful mood swings on her periods, it's like living with a monster ..and then they say 'but that's normal isn't it, she is 14?'... so it's hard to know what to do (11;F; 20-29;F)*

## Box 5. Anxiety associated with the complexity of presentations of adolescent emotional distress

*... you feel that there are these big 'no go areas' in teenage consultations which loom over you like a black cloud (11;F; 20-29; S)*

*They are missing school, in trouble with the police, youth offending team or not uncommonly the parent just comes by themselves...I never get a 14 year old acting out saying 'you know, I'm in trouble with my mam and dad' ( 10;M;40-49;P)*

*they are in absolute crisis one minute and then you see them a week later and they can hardly remember what it was all about (07;M; 40-49;P)*

*Its always a worry isn't it that you just completely get it wrong..I mean I'm conscious of this. I had someone in on Monday, parents, whose son had just hung himself at 21. I'd never seen him, he was a patient here. They had no idea anything was wrong. Nobody did....there is always that underlying things isn't there, that you might miss something catastrophic.. (01;F;50-59;S)*

## Anxiety Box iii

## Box 5. Anxiety related to interacting with young people

*there's no such thing as a typical teenage presentation ( 13;F; 30-29;S)*

*...and her mum said, as mum's do, oh she has dreadful mood swings on her periods, it's like living with a monster ..and then they say 'but that's normal isn't it, she is 14?'... so it's hard to know what to do (11;F; 20-29;F)*

## Box 6. Anxiety associated with the complexity of presentations of adolescent emotional distress

*... you feel that there are these big 'no go areas' in teenage consultations which loom over you like a black cloud (11;F; 20-29; S)*

*They are missing school, in trouble with the police, youth offending team or not uncommonly the parent just comes by themselves...I never get a 14 year old acting out saying 'you know, I'm in trouble with my mam and dad' ( 10;M;40-49;P)*

*they are in absolute crisis one minute and then you see them a week later and they can hardly remember what it was all about (07;M; 40-49;P)*

*Its always a worry isn't it that you just completely get it wrong..I mean I'm conscious of this. I had someone in on Monday, parents, whose son had just hung himself at 21. I'd never seen him, he was a patient here. They had no idea anything was wrong. Nobody did....there is always that underlying things isn't there, that you might miss something catastrophic.. (01;F;50-59;S)*

*Uncertainty is very key to this group when you're looking - in terms of depression and suicide risk and things like that, you know, it's standard. Young people particularly young males are quite at risk of just going off and doing something. (04; F; 40-49;S)*

*The main anxiety is what to do. (07;M; 40-49;P)*

1  
2  
3  
4  
5  
6  
7 Contributorship: The data presented here represents the open codes analysis which  
8 was led by JR with input from AC and JF. Further analysis of the axial and selective  
9 codes is presented elsewhere.

10  
11 **Acknowledgments, Competing interests, Funding.**

12  
13  
14 **Acknowledgements.** The authors wish to thank the GP participants for their time  
15 and candour, and their practises for supporting the research; and the RCGP  
16 Scientific Foundation Board which awarded a grant to cover the transcription costs.  
17


18  
19 **Competing interests.** We declare no competing interests.  
20

21 **Funding:** the RCGP Scientific Board awarded a grant to cover the transcription  
22 costs. SFB-2008-06  
23

24  
25 **Ethical approval:** Hull & East Riding Local Ethics Committee. REC Reference No:  
26 08/H1304/97.  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

For peer review only



## Reference list

1. Green H, McGinnity A, Meltzer H, Ford T, Goodman R. Mental Health of Children and Young People in Great Britain, 2004 : summary report. London: Office of National Statistics, 2005.
2. Singleton N, Bumpstead R, O'Brien M, Lee A, Meltzer H. Psychiatric morbidity in adults living in private households, 2000. Norwich: The Stationery Office, 2001.
3. Hawton K, Saunders KE, O'Connor RC. Self-harm and suicide in adolescents. *Lancet* 2012;379(9834):2373-82.
4. Kramer T, Garralda M. Child and adolescent mental health problems in primary care. *Advances in Psychiatric Treatment* 2000;6:287-94.
5. Hickie I, Fogarty A, Davenport T, Luscombe G, Burns J. Responding to experiences of young people with common mental health problems attending Australian general practice. *Medical Journal of Australia* 2007;187:S47-S52.
6. Gledhill J, Garralda M. The short-term outcome of depressive disorder in adolescents attending primary care: a cohort study. *Soc Psychiatry Psychiatr Epidemiol* 2011;46(10):993-1002.
7. Martinez R, Reynolds S, Howe A. Factors that influence the detection of psychological problems in adolescents attending general practices. *Br J Gen Pract* 2006;56(529):594-9.
8. Haller D, Sancu L, Sawyer S, Patton G. The identification of young people's emotional distress: a study in primary care. *Br J Gen Pract* 2009;59(560):e61-70.
9. Mauerhofer A, Berchtold A, Michaud PA, Suris JC. GPs' role in the detection of psychological problems of young people: a population-based study. *Br J Gen Pract* 2009;59(566):e308-14.
10. Churchill R. Child and Adolescent Mental Health. In: Cohen A, editor. *Delivering mental health for primary care: an evidence based approach*. London: RCGP, 2008:157-84.
11. Iliffe S, Gledhill J, da Cunha F, Kramer T, Garralda E. The recognition of adolescent depression in general practice: issues in the acquisition of new skills. *Primary Care Psychiatry* 2004;9(2):51-56.
12. Goldberg D, Huxley P. *Common mental disorders: a bio-social model*. London: Tavistock/Routledge, 1992.
13. Pilling S, Whittington C, Taylor C, Kendrick T. Identification and care pathways for common mental health disorders: summary of NICE guidance. *BMJ* 2011;342:d2868.
14. Dowrick C. *Beyond Depression: a new approach to understanding and management*. 2nd ed. Oxford: Oxford University Press, 2009.
15. Reeve J. Interpretive medicine: Supporting generalism in a changing primary care world. *Occas Pap R Coll Gen Pract* 2010(88):1-20, v.
16. Dowrick C, May C, Richardson M, Bundred P. The biopsychosocial model of general practice: rhetoric or reality? *Br J Gen Pract* 1996;46(403):105-7.
17. Karasz A, Dowrick C, Byng R, Buszewicz M, Ferri L, Hartman TCO, et al. What we talk about when we talk about depression: doctor-patient conversations and treatment decision outcomes. *Br J Gen Pract* 2012;62(594):e55-e63.
18. Kennedy I. Getting it right for children and young people: overcoming cultural barriers in the NHS so as to meet their needs. A review by Sir Ian Kennedy. London: Department of Health, 2010.
19. Layard R. How mental health loses out in the NHS. London: London School of Economics and Political Science, 2012.
20. Department for Education and Skills. *Every Child Matters: change for children* London: HMSO, 2002.
21. Department of Health. *Child and Adolescent Mental Health-National Service Framework for Children, Young People and Maternity Services* London: Department of Health 2004.
22. Department of Health. *Healthy Lives, Brighter Futures* London: The Stationery Office, 2009.

1. Green H, McGinnity A, Meltzer H, Ford T, Goodman R. *Mental Health of Children and Young People in Great Britain, 2004 : summary report*. London: Office of National Statistics, 2005. Formatted: Font: Calibri, 11 pt
2. Singleton N, Bumpstead R, O'Brien M, Lee A, Meltzer H. *Psychiatric morbidity in adults living in private households, 2000*. Norwich: The Stationery Office, 2001.
3. Hawton K, Saunders KE, O'Connor RC. Self-harm and suicide in adolescents. *Lancet* 2012;379(9834):2373-82. Formatted: Font: Calibri, 11 pt, Italic
4. Kramer T, Garralda M. Child and adolescent mental health problems in primary care. *Advances in Psychiatric Treatment* 2000;6:287-94. Formatted: Font: Calibri, 11 pt
5. Hickie I, Fogarty A, Davenport T, Luscombe G, Burns J. Responding to experiences of young people with common mental health problems attending Australian general practice. *Medical Journal of Australia* 2007;187:S47-S52. Formatted: Font: Calibri, 11 pt, Italic
6. Gledhill J, Garralda M. The short-term outcome of depressive disorder in adolescents attending primary care: a cohort study. *Soc Psychiatry Psychiatr Epidemiol* 2011;46(10):993-1002. Formatted: Font: Calibri, 11 pt, Italic
7. Martinez R, Reynolds S, Howe A. Factors that influence the detection of psychological problems in adolescents attending general practices. *Br J Gen Pract* 2006;56(529):594-9. Formatted: Font: Calibri, 11 pt
8. Haller D, Sancu L, Sawyer S, Patton G. The identification of young people's emotional distress: a study in primary care. *Br J Gen Pract* 2009;59(560):e61-70. Formatted: Font: Calibri, 11 pt, Italic
9. Mauerhofer A, Berchtold A, Michaud PA, Suris JC. GPs' role in the detection of psychological problems of young people: a population-based study. *Br J Gen Pract* 2009;59(566):e308-14. Formatted: Font: Calibri, 11 pt
10. Churchill R. *Child and Adolescent Mental Health*. In: Cohen A, editor. *Delivering mental health for primary care: an evidence based approach*. London: RCGP, 2008:157-84. Formatted: Font: Calibri, 11 pt, Italic
11. Iliffe S, Gledhill J, da Cunha F, Kramer T, Garralda E. The recognition of adolescent depression in general practice: issues in the acquisition of new skills. *Primary Care Psychiatry* 2004;9(2):51-56. Formatted: Font: Calibri, 11 pt
12. Goldberg D, Huxley P. *Common mental disorders: a bio-social model*. London: Tavistock/Routledge, 1992. Formatted: Font: Calibri, 11 pt, Italic
13. Dowrick C. *Beyond Depression: a new approach to understanding and management*, 2nd ed. Oxford: Oxford University Press, 2009. Formatted: Font: Calibri, 11 pt
14. Reeve J. Interpretive medicine: Supporting generalism in a changing primary care world. *Occas Pap R Coll Gen Pract* 2010(88):1-20, v. Formatted: Font: Calibri, 11 pt, Italic
15. Dowrick C. Reasons to be cheerful? Reflections on GPs' responses to depression. *Br J Gen Pract* 2009;59(566):636-7. Formatted: Font: Calibri, 11 pt
16. Dowrick C, May C, Richardson M, Bundred P. The biopsychosocial model of general practice: rhetoric or reality? *Br J Gen Pract* 1996;46(403):105-7. Formatted: Font: Calibri, 11 pt, Italic
17. Karasz A, Dowrick C, Byng R, Buszewicz M, Ferri L, Hartman TCO, et al. What we talk about when we talk about depression: doctor-patient conversations and treatment decision outcomes. *Br J Gen Pract* 2012;62(594):e55-e63. Formatted: Font: Calibri, 11 pt
18. Kennedy I. Getting it right for children and young people: overcoming cultural barriers in the NHS so as to meet their needs. A review by Sir Ian Kennedy. London: Department of Health, 2010. Formatted: Font: Calibri, 11 pt, Italic
19. Layard R. *How mental health loses out in the NHS*. London: London School of Economics and Political Science, 2012. Formatted: Font: Calibri, 11 pt
20. Department for Education and Skills. *Every Child Matters: change for children* London: HMSO, 2002.
21. Department of Health. *Child and Adolescent Mental Health-National Service Framework for Children, Young People and Maternity Services* London: Department of Health 2004. Formatted: Font: Calibri, 11 pt, Italic
22. Department of Health. *Healthy Lives, Brighter Futures* London: The Stationery Office, 2009. Formatted: Font: Calibri, 11 pt
23. National Institute for Health & Clinical Excellence. *Depression in children and young people: identification and management in primary, community and secondary care*. *National Clinical Practice Guidelines*. London: NICE, 2005. Formatted: Font: Calibri, 11 pt, Italic



24. Tait L. To disclose or not to disclose psychological problems to GPs. *Br J Gen Pract* 2009;59(566):638-9.
25. Biddle L, Donovan JL, Gunnell D, Sharp D. Young adults' perceptions of GPs as a help source for mental distress: a qualitative study. *Br J Gen Pract* 2006;56(533):924-31.
26. Glaser B, Strauss A. *The discovery of grounded theory*. Chicago: Aldine, 1967.
27. Charmaz K. *Constructing Grounded Theory: A practical guide through qualitative analysis*. Thousand Oaks, California: Sage Publications, Inc, 2006.
28. Clarke A. *Situational analysis: Grounded Theory after the postmodern turn*. Sage Publications, Inc. Thousand Oaks, California: Sage Publications, Inc, 2005.
29. Chew-Graham CA, May CR, Perry MS. Qualitative research and the problem of judgement: lessons from interviewing fellow professionals. *Family Practice* 2002;19(3):285-89.
30. Heath I. Uncertain clarity: contradiction, meaning, and hope. *Brit J Gen Pract* 1999;49(445):651-57.
31. Schon D. *The reflective practitioner*. San Francisco: Jossey-Bass, 1983.
32. Iliffe S, Williams G, Fernandez V, Vila M, Kramer T, Gledhill J, et al. Treading a fine line: is diagnosing depression in young people just medicalising moodiness? *Br J Gen Pract* 2009;59(560):156-7.
33. Iliffe S, Gallant C, Kramer T, Gledhill J, Bye A, Fernandez V, et al. Therapeutic identification of depression in young people: lessons from the introduction of a new technique in general practice. *Br J Gen Pract* 2012;62(596):e174-82.
34. Sawyer SM, Afifi RA, Bearinger LH, Blakemore SJ, Dick B, Ezech AC, et al. Adolescence: a foundation for future health. *Lancet* 2012;379(9826):1630-40.
23. National Institute for Health & Clinical Excellence. Depression in children and young people: identification and management in primary, community and secondary care. *National Clinical Practice Guidelines*. London: NICE, 2005.
24. Tait L. To disclose or not to disclose psychological problems to GPs. *Br J Gen Pract* 2009;59(566):638-9.
25. Biddle L, Donovan JL, Gunnell D, Sharp D. Young adults' perceptions of GPs as a help source for mental distress: a qualitative study. *Br J Gen Pract* 2006;56(533):924-31.
26. Glaser B, Strauss A. *The discovery of grounded theory*. Chicago: Aldine, 1967.
27. Charmaz K. *Constructing Grounded Theory: A practical guide through qualitative analysis*. Thousand Oaks, California: Sage Publications, Inc, 2006.
28. Clarke A. *Situational analysis: Grounded Theory after the postmodern turn*. Sage Publications, Inc. Thousand Oaks, California: Sage Publications, Inc, 2005.
29. Chew-Graham CA, May CR, Perry MS. Qualitative research and the problem of judgement: lessons from interviewing fellow professionals. *Family Practice* 2002;19(3):285-89.
30. Heath I. Uncertain clarity: contradiction, meaning, and hope. *Brit J Gen Pract* 1999;49(445):651-57.
31. Schon D. *The reflective practitioner*. San Francisco: Jossey-Bass, 1983.
32. Dowrick C. Reasons to be cheerful? Reflections on GPs' responses to depression. *Br J Gen Pract* 2009;59(566):636-7.
33. Iliffe S, Williams G, Fernandez V, Vila M, Kramer T, Gledhill J, et al. Treading a fine line: is diagnosing depression in young people just medicalising moodiness? *Br J Gen Pract* 2009;59(560):156-7.

Formatted: Font: Calibri, 11 pt, Italic

Formatted: Font: Calibri, 11 pt

Formatted: Font: Calibri, 11 pt, Italic

Formatted: Font: Calibri, 11 pt

Formatted: Font: Calibri, 11 pt, Italic

Formatted: Font: Calibri, 11 pt

Formatted: Font: Calibri, 11 pt, Italic

Formatted: Font: Calibri, 11 pt

Formatted: Font: Calibri, 11 pt, Italic

Formatted: Font: Calibri, 11 pt

Formatted: Font: Calibri, 11 pt, Italic

Formatted: Font: Calibri, 11 pt

Formatted: Font: Calibri, 11 pt, Italic

Formatted: Font: Calibri, 11 pt

Formatted: Font: Calibri, 11 pt, Italic

Formatted: Font: Calibri, 11 pt

Formatted: Font: Calibri, 11 pt, Italic

Formatted: Font: Calibri, 11 pt

Formatted: Font: Calibri, 11 pt, Italic

Formatted: Font: Calibri, 11 pt

Formatted: Font: Calibri, 11 pt, Italic

Formatted: Font: Calibri, 11 pt



1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

34. Hiffe S, Gallant C, Kramer T, Gledhill J, Bye A, Fernandez V, et al. Therapeutic identification of depression in young people: lessons from the introduction of a new technique in general practice. *Br J Gen Pract* 2012;62(596):e174-82.

35. Sawyer SM, Afifi RA, Bearinger LH, Blakemore SJ, Dick B, Ezech AC, et al. Adolescence: a foundation for future health. *Lancet* 2012;379(9826):1630-40.

For peer review only