

BMJ Open

Ultra-processed foods and added sugars in the US diet

Journal:	<i>BMJ Open</i>
Manuscript ID:	bmjopen-2015-009892
Article Type:	Research
Date Submitted by the Author:	03-Sep-2015
Complete List of Authors:	Martinez Steele, Euridice; University of Sao Paulo, Department of Nutrition, School of Public Health Baraldi, Larissa; University of Sao Paulo, Department of Nutrition, School of Public Health Louzada, Maria Laura; University of Sao Paulo, Department of Nutrition, School of Public Health; University of São Paulo, Center for Epidemiological Studies in Health and Nutrition Moubarac, Jean-Claude; University of São Paulo, Center for Epidemiological Studies in Health and Nutrition Mozaffarian, Dariush; Friedman School of Nutrition Science and Policy, Tufts University, Monteiro, Carlos; University of Sao Paulo, Department of Nutrition, School of Public Health; University of São Paulo, Center for Epidemiological Studies in Health and Nutrition
Primary Subject Heading:	Nutrition and metabolism
Secondary Subject Heading:	Public health
Keywords:	ultra-processed, added sugars, dietary intake, NHANES, US

SCHOLARONE™
Manuscripts

Ultra-processed foods and added sugars in the US diet

Names of authors:

Eurídice Martínez Steele (EMS; PHD student)^{ab},

Larissa Galastri Baraldi (LGB; PHD student)^{ab},

Maria Laura da Costa Louzada (ML; PHD student)^{ab},

Jean- Claude Moubarac (JCM, Researcher)^b,

Dariush Mozaffarian (DM; Professor)^c,

Carlos Augusto Monteiro (CAM; Professor)^{ab}

^aDepartment of Nutrition, School of Public Health, University of São Paulo, São Paulo, Brazil

^bCenter for Epidemiological Studies in Health and Nutrition, University of São Paulo, São Paulo, Brazil

^cFriedman School of Nutrition Science and Policy, Tufts University, Boston, MA

Corresponding author:

Carlos Augusto Monteiro, Departamento de Nutrição, Faculdade de Saúde Pública, Universidade de São Paulo, Av. Dr. Arnaldo, 715, São Paulo 01246-907, Brazil. E-mail: carlosam@usp.br

Keywords: ultra-processed, added sugars, dietary intake, NHANES, US.

ABSTRACT

Objectives: To investigate the contribution of ultra-processed foods to the intake of added sugars in the US.

Food items were classified according to extent and purpose of industrial food processing. Ultra-processed foods were defined as formulations manufactured mostly or entirely from substances extracted from foods or obtained from further processing of constituents of foods.

Design: Cross- sectional study.

Setting: *National Health and Nutrition Examination Survey 2009-2010.*

Participants: We evaluated 9,317 participants aged 1+ years with at least one 24-hour dietary recall.

Main outcome measures: Average dietary content of added sugars and proportion of individuals consuming more than 10% of total energy from added sugars.

Data analysis: Gaussian and Poisson regressions estimated the association between consumption of ultra-processed foods and intake of added sugars. All models incorporated survey sample weights and adjusted for age, sex, race/ethnicity, and family income.

Results: Ultra-processed foods comprised 57.5% of energy intake, and contributed 89.7% of the energy intake from added sugars. The content of added sugars in ultra-processed foods (20.8% of calories) was 9-fold higher than in processed foods (2.3%) and 5-fold higher than in unprocessed or

minimally processed foods and processed culinary ingredients grouped together (3.6%). In both unadjusted and adjusted models, each increase of 5 percentage points in proportional energy intake from ultra-processed foods increased the proportional energy intake from added sugars by 1 percentage point. Consumption of added sugars increased linearly across quintiles of ultra-processed food consumption: from 7.8% of total energy in the lowest quintile to 19.2% in the highest. A total of 85.8% of Americans in the highest quintile exceeded the recommended limit of 10% energy from added sugars, compared with 28.9% in the lowest.

Conclusions: Decreasing the consumption of ultra-processed foods could be an effective way of reducing the excessive intake of added sugars in the US.

Strengths and limitations of this study:

- Use of a large, nationally representative sample of the US population, increasing generalizability.
- Use of data on added sugars rather than total sugars or sugar-sweetened beverages, which corresponds to guidelines relevant area of prioritization.
- Unlike most articles which have focused on specific food items such as soft drinks or fast food, our study evaluates the impact of a comprehensive group of products whose consumption is increasing exponentially in most countries.
- Dietary data obtained by 24-hour recalls is subjected to potential error and bias.

- Information indicative of food processing is not consistently determined for all food items in NHANES, which could lead to modest over or underestimation of the consumption of ultra-processed foods.

For peer review only

1 INTRODUCTION

2 Increasing policy attention has focused on added sugars, including by the World
3 Health Organization (WHO)(1), the United Kingdom National Health System(2),
4 the Canadian Heart and Stroke Foundation(3), the American Heart Association
5 (AHA)(4), and the US Dietary Guidelines Advisory Committee (USDGAC)(5).

6 These reports concluded that a high intake of added sugars increases risk for
7 weight gain(1,4,5), excess body weight(5) and obesity(3,5); type 2 diabetes
8 mellitus(3,5); higher serum triglycerides(5) and high blood cholesterol(3); higher
9 blood pressure(5) and hypertension(5); stroke(3,5); coronary heart disease(3,5);
10 cancer(3); and dental caries(1,3,5). Moreover, foods higher in added sugars
11 are often a source of empty calories with minimum essential nutrients or dietary
12 fiber(6-8), which displace more nutrient-dense foods(9) and lead, in turn, to
13 simultaneously overfed and undernourished individuals.

14 All reports recommended limiting intake of added sugars(1,3-5). In the US, the
15 USDGAC recommended limiting added sugars to no more than 10% of total
16 calories. This is a challenge, as recent consumption of added sugars in the US
17 amounted to almost 15% of total calories in 2005-2010(10,11).

18 To design and implement effective measures to reduce added sugars, their
19 dietary sources must be clearly identified. Added sugars can be consumed
20 either as ingredients of dishes or drinks prepared from scratch by consumers or
21 cook, or as ingredients of food products manufactured by the food
22 industry. According to market disappearance data from 2014, more than three
23 quarters of the sugar and high fructose corn syrup available for human
24 consumption in the US were used by the food industry(12). This suggests food

1
2
3 25 products manufactured by the industry could have an important role in the
4
5 26 excess added sugars consumption in the US. However, to assess this role, it is
6
7 27 essential to consider the contribution of manufactured food products to both
8
9 28 total energy intake and the energy intake from added sugars, and, more
10
11 29 relevantly, to quantify the relationship between their consumption and the total
12
13 30 dietary content of added sugars. To address these questions, we performed an
14
15 31 investigation utilizing 2009-2010 *National Health and Nutrition Examination*
16
17 32 *Survey (NHANES)*.
18
19
20
21
22 33
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

34 **SUBJECTS AND METHODS**

35 **Data source, population and sampling**

36 We utilized nationally representative data from the 2009-2010 *National Health*
37 *and Nutrition Examination Survey (NHANES)*, specifically the dietary
38 component *What we eat in America (WWEIA)*(13).

39 NHANES is a continuous, nationally representative, cross-sectional survey of
40 the non-institutionalized, civilian US residents(14). NHANES sample was
41 obtained by using a complex, stratified, multi-stage probability cluster sampling
42 design, based on the selection of counties, blocks, households, and the number
43 of people within households(14). In order to improve the estimate precision and
44 reliability, NHANES 2009-2010 oversampled the following subgroups: Hispanic,
45 Non-Hispanic black, Non-Hispanic white and Other persons at or below 130%
46 of the federal poverty level and Non-Hispanic white and Other persons aged 80
47 + years(14).

48 All NHANES examinees were eligible for two 24-hour dietary recall interviews.
49 The first dietary recall interview was collected in-person in the mobile
50 examination center (MEC)(15) while the second was collected by telephone 3 to
51 10 days later(16). Dietary interviews were conducted by trained interviewers
52 using the validated(17-19) *US Department of Agriculture Automated Multiple-*
53 *Pass Method (AMPM)*(20).

54 Among the 13,272 people screened in NHANES 2009-2010, 10,537 (79.4%)
55 participated in the household interview and 10253 (77.3%) also participated in
56 the MEC health examination(21). Of these, 9,754 individuals provided one day
57 of complete dietary intakes and 8,406 provided two days(22).

We evaluated 9,317 survey participants aged 1 year and above who had one day 24-hour dietary recall data and had not been breast-fed on either of the two days. These individuals had similar socio-demographic characteristics to the full sample of 10,109 participants interviewed.

Food classification according to processing

We classified all recorded food items (N=280,132 Food Codes) according to NOVA, a food classification based on the extent and purpose of industrial food processing(23-25). This classification includes 4 groups: “unprocessed or minimally processed foods” (such as fresh, dry or frozen fruits or vegetables, grains, legumes, meat, fish and milk); “processed culinary ingredients” (including table sugar, oils, fats, salt, and other substances extracted from foods or from nature, and used in kitchens to make culinary preparations), “processed foods” (foods manufactured with the addition of salt or sugar or other substances of culinary use to unprocessed or minimally processed foods, such as simple cheese, bread and canned food), and “ultra-processed foods” (formulations manufactured mostly or entirely from starches, sugars, oils, fats, proteins, and other substances extracted from foods or obtained from the further processing of constituents of foods or other organic sources, with little if any whole foods). A detailed definition of each food group and examples of food items classified in each group are shown in **Supplementary Table 1**. The rationale underlying the classification is described elsewhere(26-29).

For all food items (Food Codes) judged to be a handmade recipe, the classification was applied to the underlying ingredients (Standard Reference Codes -SR Codes-) obtained from the USDA Food and Nutrient Database for

82 Dietary Studies (FNDDS) 5.0(30). Refer to Online Supplementary Material
83 (OSM) for further details.

84 **Assessing energy and added sugar contents**

85 For this study, we used Food Code energy values as provided by NHANES.

86 For handmade recipes, we calculated the underlying ingredient (SR Code)
87 energy values using variables from both *FNDDS 5.0(30)* and *USDA National*
88 *Nutrient Database for Standard Reference, Release 24 (SR24)(31)*. Refer to
89 OSM for further details.

90 Data on added sugars per Food Code and per SR Code were obtained by
91 merging the Food Patterns Equivalents Database (FPED) 2009-2010 and the
92 Food Patterns Equivalents Ingredients Database (FPID) 2009-2010(32). Added
93 sugars are defined in these databases as “sugars that are added to foods as an
94 ingredient during preparation, processing, or at the table. Added sugars do not
95 include naturally occurring sugars (e.g., lactose in milk, fructose in fruits).

96 Examples of added sugars include brown sugar, cane sugar, confectioners’
97 sugar, granulated sugar, dextrose, white sugar, corn syrup and corn syrup
98 solids, molasses, honey, and all types of syrups such as maple syrup, table
99 syrups, and pancake syrup”(32). These two databases express the content of
100 added sugars in teaspoons per 100 g. Teaspoons were converted into grams
101 using the factor 4.2 g/teaspoon and into kcal using the factor 3.87 kcal/g.

102 **Data Analysis**

103 We utilized all available dietary data for each participant, using means of both
104 recall days when available (86% of participants). Food items were sorted into

105 mutually exclusive food subgroups within Unprocessed or minimally processed
106 foods (n= 11), Processed culinary ingredients (n=4), Processed foods (n=4) and
107 Ultra-processed foods (n=18), as shown in Table 1. First, we evaluated the
108 contributions of each of the NOVA food groups and subgroups to total energy
109 and to the energy from added sugars. Next, we calculated the average content
110 of added sugars in the overall US diet and in fractions of this diet composed by
111 each of the NOVA food groups and subgroups. We also calculated the dietary
112 content of added sugars in the group of unprocessed or minimally processed
113 foods combined with the group of processed culinary ingredients, as foods
114 belonging to these two groups are usually combined together in culinary
115 preparations and therefore consumed together.

116 We used Gaussian regression to estimate the association between the dietary
117 contribution of ultra-processed foods and the dietary content of added sugars,
118 each expressed as proportions of total energy. This association was also
119 explored after adjusting for the proportion of added sugars in non-ultra-
120 processed energy intake. Dietary contribution of ultra-processed foods was
121 transformed using restricted cubic spline functions to allow for nonlinearity.

122 The average content of added sugars in the overall diet was compared across
123 quintiles of the dietary contribution of ultra-processed foods. Poisson
124 regression was used to assess whether the percentage of diets with more than
125 10% or 20% of total energy from added sugars increased across quintiles. This
126 increase was also evaluated across demographic subgroups in stratified
127 analysis. Tests of linear trend were performed in order to evaluate the effect of
128 quintiles as a single continuous variable.

129 All regression models were adjusted for age (1-5 years, 6-11 years, 12-19
130 years, 20–39 years, 40–59 years, 60 + years), sex, race/ethnicity (Mexican-
131 American, Other Hispanic, Non-Hispanic White, Non-Hispanic Black, Other
132 Race including Multi-Racial) and ratio of family income to poverty (categorized
133 based on Supplemental Nutrition Assistance Program (SNAP) eligibility as
134 0.00–1.30, >1.30–3.50, and >3.50 and above)(14). As 833 participants had
135 missing values on income to poverty, adjusted analysis included 8,484
136 individuals.

137 NHANES survey sample weights were used in all analyses to account for
138 differential probabilities of selection for the individual domains, nonresponse to
139 survey instruments, and differences between the final sample and the total US
140 population. The Taylor series linearization variance approximation procedure
141 was used for variance estimation in all analysis in order to account for the
142 complex sample design and the sample weights(14).

143 To minimize chance findings from multiple comparisons, statistical hypotheses
144 were tested using a two-tailed $p<0.001$ level of significance. Data were
145 analyzed using Stata statistical software package version 12.1.

146 RESULTS

147 Distribution of total energy intake by food groups

148 The average US daily energy intake in 2009-2010 was 2044.6 kcal, and nearly
149 3 in 5 calories (57.5%) came from ultra-processed foods (**Table 1**).

150 Unprocessed or minimally processed foods contributed 30.3% of total calories,
151 processed foods an additional 9.3%, and processed culinary ingredients the
152 remaining 2.9%. The most common ultra-processed foods in terms of energy
153 contribution were breads; soft drinks, fruit drinks, and milk drinks; cakes,
154 cookies, and pies; salty-snacks; frozen and shelf-stable plates; pizza; and
155 breakfast cereals. Meat, fruit, and milk provided the most calories among
156 unprocessed or minimally processed foods; ham and cheese, the most calories
157 among processed foods; and table sugar and plant oils, the most calories
158 among processed culinary ingredients.

159

Table 1. Distribution of the total energy intake and of the energy intake from added sugars according to food groups, and the mean content of added sugars of each food group. US population aged 1 + years (NHANES 2009-2010) (N=9,317)

Food groups	Mean energy intake		Mean energy intake from added sugars		Mean content of added sugars
	Absolute (kcal/day)	Relative (% of total energy intake)	Absolute (kcal/day)	Relative (% of total energy intake from added sugars)	% of energy from added sugars
Unprocessed or minimally processed foods	591.1	30.3	0.0	0.0	0.0
Meat (includes poultry)	170.3	8.3	0.0	0.0	0.0
Fruit ¹	96.8	5.2	0.0	0.0	0.0
Milk and plain yoghurt	96.8	5.1	0.0	0.0	0.0
Grains	54.5	2.9	0.0	0.0	0.0
Roots and tubers	32.2	1.6	0.0	0.0	0.0
Eggs	29.1	1.5	0.0	0.0	0.0
Pasta	29.3	1.4	0.0	0.0	0.0
Legumes	16.7	0.8	0.0	0.0	0.0
Fish and sea food	15.7	0.8	0.0	0.0	0.0
Vegetables	13.9	0.8	0.0	0.0	0.0
Other unprocessed or minimally processed foods ²	35.6	1.7	0.0	0.0	0.0
Processed culinary ingredients	62.9	2.9	23.9	8.7	38.3
Table sugar ³	24.2	1.1	23.9	8.7	98.4

Plant oils	26.8	1.2	0.0	0.0	0.0
Animal fats ⁴	10.9	0.5	0.0	0.0	0.0
Other processed culinary ingredients ⁵	0.9	0.04	0.0	0.0	0.0
Unprocessed or minimally processed foods + Processed culinary ingredients	653.9	33.2	23.9	8.7	3.6
Processed foods	203.2	9.3	2.4	1.6	2.3
Cheese	77.7	3.7	0.0	0.0	0.0
Ham and other salted, smoked or canned meat or fish	26.2	1.3	0.3	0.2	1.4
Vegetables and other plant foods preserved in brine	13.3	0.7	1.5	0.9	13.5
Other processed foods ⁶	86	3.7	0.6	0.4	1.3
Ultra-processed foods	1187.4	57.5	254.5	89.7	20.8
Breads ⁷	190.5	9.6	10.4	7.8	5.6
Cakes, cookies and pies	117.2	5.5	28	10.9	23.9
Salty-snacks	90.7	4.4	1.2	0.8	1.4
Frozen and shelf-stable plate meals	79.7	3.9	1.1	0.8	1.7
Soft drinks, carbonated	77.2	3.6	70.9	16.5	69.6
Pizza (ready-to-eat/heat)	77.2	3.3	2.3	1.4	2.9
Fruit drinks ⁸	65	3.2	51.9	13.5	67.3
Breakfast cereals	56.1	3	13.7	7.1	23.5

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49

Sauces, dressings and gravies	52.4	2.5	4.5	3	9.8
Reconstituted meat or fish products	49.6	2.3	0.7	0.6	1.9
Ice cream and ice pops	49.3	2.3	18.2	6.1	36.9
Sweet-snacks	48.3	2.3	17.9	6.7	37.8
Milk drinks ⁹	35.7	1.9	11.1	4.8	33.3
Desserts ¹⁰	35.7	1.8	17.8	7.3	48.1
French fries and other potatoe products ¹¹	37.7	1.7	0.0	0.0	0.0
Sandwiches and hamburgers on bun (ready-to-eat/heat)	31.9	1.4	1.3	0.6	4.3
Instant and canned soups	15.1	0.9	0.2	0.1	0.8
Other ultra-processed foods ¹²	78.2	3.7	3	1.5	7.9
Total	2044.6	100.0	280.8	100.0	13.4

- 162 1Including freshly squeezed juices
- 163 2Including nuts and seeds (unsalted); yeast; dried fruits (without added sugars) and vegetables; non pre-sweetened, non-whitened, non-
164 flavored coffee and tea; coconut water and meat; homemade soup and sauces (with no underlying ingredients); flours; tapioca
- 165 3Including honey, molasses, maple syrup (100%)
- 166 4Including unsalted butter, lard and cream
- 167 5Including starches; coconut and milk cream; unsweetened baking chocolate, cocoa powder and gelatin powder; vinegar; baking powder and
168 baking soda
- 169 6Including salted or sugared nuts and seeds; peanut, sesame, cashew and almond butter or spread; beer and wine

- 170 7Including all types of bread. Processed bread made of flour, water, salt, leavening agents and possibly walnuts, dried fruits and other whole
171 foods, were included under this group as well, because of the low consumption
- 172 8Including fruit and fruit-flavored, non-carbonated and other sweetened drinks, including presweetened tea and coffee, energy drinks, sports
173 drinks with no milk added, nonalcoholic wine
- 174 9Including flavored yogurt sweetened with sugar or with low-calorie sweetener, milk-shake, soymilk
- 175 10Including ready-to-eat and dry-mix desserts such as pudding; sugar based ingredients such as whipped cream; sweetened canned fruit and
176 fruit sauce
- 177 11Including hash browns, potato puffs, stuffed potatoes, onion rings (ready-to-eat/heat)
- 178 12Including soy products such as meatless patties and fish sticks; babyfood and baby formula; dips, spreads, mustard and catsup; salted butter
179 and margarine; sugar substitutes, sweeteners and all syrups (excluding 100% maple syrup); distilled alcoholic drinks

180 **Distribution of energy intake from added sugars by food groups**

181 The average US daily intake of added sugars was 280.8 kcal (Table 1).
182 Notably, almost 90% of this (89.7%) came from ultra-processed foods. The
183 main sources of added sugars among ultra-processed foods were: soft drinks
184 (16.5% of US intake of added sugars), fruit drinks (13.5%), milk drinks (4.8%);
185 cakes, cookies, and pies (10.9%); breads (7.8%); desserts (7.3%); breakfast
186 cereals (7.1%); sweet snacks (6.7%), and ice creams and ice pops (6.1%). In
187 contrast, only 8.7% of the added sugars in the US diet came from processed
188 culinary ingredients (table sugar consumed as part of dishes or drinks prepared
189 from scratch by consumers or cook), and only 1.6% from processed foods.

190 The average content of added sugars in ultra-processed foods (20.8% of
191 calories) was 9-fold higher than in processed foods (2.3%) and 5-fold higher
192 than in unprocessed or minimally processed foods and processed culinary
193 ingredients grouped together (3.6%) (Table 1).

194 **Association between consumption of ultra-processed foods and added**
195 **sugar intake**

196 In both unadjusted and multivariable-adjusted restricted cubic splines Gaussian
197 regression analyses, a strong linear association was identified between the
198 dietary contribution (percentage of calories) of ultra-processed foods and the
199 dietary content (percentage of calories) in added sugars (unadjusted coefficient
200 for linear term=0.21, 95% CI: 0.18 to 0.24) (**Figure 1**).

201 **Figure 1.**

202 There was little evidence of nonlinearity in the restricted cubic spline model
203 (Wald test for linear term $p < 0.0001$; Wald test for all non-linear terms $p = 0.19$ –
204 unadjusted model-). The strength of the association remained fairly the same
205 after adjusting for the proportion of added sugars in non-ultra-processed energy
206 intake (coefficient for linear term = 0.19, 95% CI: 0.17 to 0.23) and for age, sex,
207 race/ethnicity, and family income (coefficient for linear term = 0.22, 95% CI: 0.18
208 to 0.26).

209 Overall, each increase in 5 percentage points of energy in consumption of ultra-
210 processed foods was associated with 1 higher percentage point of energy in the
211 consumption of added sugars.

212 Across quintiles of energy-adjusted ultra-processed food consumption, the
213 intake of added sugars increased substantially and monotonically, from 7.8% of
214 total calories in the lowest quintile to 19.2% in the highest. Across the same
215 quintiles, the proportion of individuals consuming more than 10% of total energy
216 from added sugars (61.5% in the total population) increased from 28.9% to
217 85.8%, respectively. An even more pronounced increase was seen in the
218 proportion of individuals consuming more than 20% of their total energy from
219 added sugars: from 3.6% in the lowest quintile to 41.1% in the highest (**Table**
220 **2**). Similar increases were seen in stratified analysis by major demographic
221 subgroups (**Supplementary Table 2**). The magnitude and the statistical
222 significance of the association between the dietary contribution of ultra-
223 processed foods and the dietary content in added sugars did not change with
224 adjustment for sex, age, race/ethnicity, and family income.

Table 2. Indicators of the dietary content in added sugars according to the dietary contribution of ultra-processed foods. US population aged 1 + years (NHANES 2009-2010)

Dietary contribution of ultra-processed foods (% of total energy intake)		Indicators						
		% of total energy intake from added sugars	Participants with more than 10% of total energy intake from added sugars			Participants with more than 20% of total energy intake from added sugars		
Quintiles	Mean (range)	Mean	%	PR ¹	PRadj ²	%	PR ¹	PRadj ²
1st (n=1,955)	32.5 (0 to 42.7)	7.8	28.9	1	1	3.6	1	1
2nd (n=1,901)	48.6 (42.7 to 53.9)	11.2	50.1	1.7	1.8	8.5	2.4	2.5
3rd (n=1,774)	58.3 (53.9 to 62.6)	13.3	65.1	2.2	2.3	15.2	4.3	4.3
4th (n=1,784)	67.2 (62.6 to 72.1)	15.7	77.7	2.7	2.7	23.9	6.7	6.5
5th (n=1,903)	80.7 (72.1 to 100)	19.2*	85.8	2.9*	2.9*	41.1	11.5*	10.9*
Total (n=9,317)	57.5 (0 to 100.0)	13.4	61.5			18.5		

*Significant linear trend across all quintiles (p<=0.001), both in unadjusted and models adjusted for sex, age group (1-5, 6–11, 12–19, 20–39, 40–59, 60 + years), race/ethnicity (Mexican-American, Other Hispanic, Non-Hispanic White, Non-Hispanic Black and Other Race - Including Multi-Racial-), and ratio of family income to poverty (SNAP 0.00–1.30, >1.30–3.50, and >3.50 and over).

1PR=Prevalence ratios estimated using Poisson regression (N=9,317)

2PRadj=Prevalence ratios adjusted for sex, age groups, race/ethnicity and ratio of family income to poverty, as above (N=8,484)

DISCUSSION

In this analysis of nationally representative data, we confirmed the excessive consumption of added sugars in the US(10,11). We also provide new evidence that ultra-processed foods represent more than half of all calories in the US diet, and contribute nearly 90% of all added sugars. Added sugars represented 1 of every 5 calories in the average ultra-processed food (20.8%), far higher than the content of added sugars in processed foods (2.3%) and in unprocessed or minimally processed foods, and processed culinary ingredients grouped together (3.6%). A strong linear relationship was found between the dietary contribution of ultra-processed foods and the dietary content of added sugars. Moreover, the risk of exceeding the recommended upper limit of 10% energy from added sugars was far higher when ultra-processed food consumption was high, and risk differences were even more pronounced for exceeding a limit of 20% energy. Notably, only those Americans in the lowest quintile of ultra-processed food consumption met the recommended limit of <10% energy from added sugars. To our knowledge, this is the first study to assess the consumption of ultra-processed foods and establish its relationship with excessive added sugar intake in the US.

The high consumption of added sugars in the US is likely contributing to excess obesity, type 2 diabetes, dyslipidemia, hypertension and coronary heart disease(1,3-5). Consequently, most dietary guidelines now recommend limiting added sugar consumption. However, such guidelines are not always clear on how to put this recommendation into practice. Our study suggests that in the US, limiting the consumption of ultra-processed foods may be a highly effective way to decrease added sugars. A reduction in ultra-processed foods should

257 also increase the intake of more healthful, minimally processed foods such as
258 milk, fruits, and nuts, and freshly-prepared dishes based on whole grains and
259 vegetables, which would produce additional health benefits beyond the
260 reduction in added sugar. Consistent with this approach, in Brazil, where the
261 consumption of added sugars is as high as in the US(33), the new dietary
262 guidelines launched in 2014 emphasize the importance of not replacing
263 unprocessed or minimally processed foods and freshly prepared dishes by
264 ultra-processed foods(34).

265 Few studies have assessed the impact of levels of food processing on the
266 nutrient profile of the US diet. One analysis using data from NHANES 2003-
267 2008(35) used a food classification system(36) including “Mixtures of combined
268 Ingredients” and “Ready-to-eat”, which are mostly ultra-processed foods and
269 together, contributed to about half of total energy intake and three-quarters of
270 energy intake from added sugars. Another study evaluated household
271 barcoded purchasing data from 2000-2012 using a classification system guided
272 by the one used in our study(37). In 2012, the mean per capita purchase of
273 “highly processed foods”, a category similar to ultra-processed foods,
274 corresponded to 61.0% of all calories and had higher adjusted median total
275 sugar content than “less processed foods”. This report did not evaluate added
276 sugars nor the contribution of processed foods to sugar intake. It also did not
277 capture non-barcoded items such as unpackaged fresh fruit, vegetables and
278 meat, or highly processed foods such as ready-to-eat store-prepared items. An
279 investigation in Canada, using 2001 household purchasing data, found that
280 ultra-processed foods are high in free sugars and that only households in the
281 lowest quintile of ultra-processed food purchasing might have met the

recommended limit of <10% energy from free sugars (9.2%)(38). Being based on household purchasing data, these two prior studies and others based on the NOVA classification system(23, 39-42) could not evaluate fraction of wasted food nor purchases at restaurants, which represent a substantial proportion of US calories. Our findings build upon and considerably extend these prior reports by evaluating food processing and added sugar intake using contemporary, nationally representative dietary intake data in the US.

Our study has several strengths. We studied a large, nationally representative sample of the US population, increasing generalizability. Use of data on added sugars rather than total sugars or sugar-sweetened beverages, corresponds to the relevant area of prioritization of recent national and international guidelines. Our investigation was based on individual consumption data, rather than market disappearance or household purchasing data which cannot account for differences between amounts purchased and amounts actually consumed.

Potential limitations should be considered. As with most population measures, dietary data obtained by 24-hour recalls is imperfect. However, the standardized methods and approach of NHANES and use of two recalls per person minimize potential error and bias, particularly for assessing population averages as focused upon in the present study. Previous studies suggest that people with obesity may underreport consumption of foods with caloric sweeteners(43) such as desserts and sweet baked goods(44, 45). If so, these biases may lead to an underestimation of the dietary contribution of ultra-processed foods and the overall intake of added sugars, but should have much less effect on the association between these. Although NHANES collects some information indicative of food processing (i.e. place of meals, product brands),

307 these data are not consistently determined for all food items, which could lead
308 to modest over or underestimation of the consumption of ultra-processed foods.

309 In conclusion, we found that ultra-processed foods contribute almost 60% of
310 calories and 90% of added sugars consumed in the US. Only Americans in the
311 lowest quintile of ultra-processed food consumption met the recommended
312 guidelines for intake of added sugars. Decreasing the consumption of ultra-
313 processed foods could be an effective way of reducing the excessive intake of
314 added sugars in the US.

315

316

317 Competing Interests.

318 "All authors have completed the ICMJE uniform disclosure form at
319 www.icmje.org/coi_disclosure.pdf and declare: no support from any
320 organisation for the submitted work; DM reports ad hoc honoraria or consulting
321 from Bunge, Haas Avocado Board, Nutrition Impact, Amarin, Astra Zeneca,
322 Boston Heart Diagnostics, GOED, and Life Sciences Research Organization;
323 and scientific advisory boards, Unilever North America and Elysium Health; no
324 other relationships or activities that could appear to have influenced the
325 submitted work."

326 Contributorship statement.

327 CAM, EMS, DM designed research; EMS, LB, ML data management; EMS, J-
328 CM, ML analyzed data; EMS, DM, CAM wrote paper; CAM, EMS had primary
329 responsibility for final content. All authors read and approved the final
330 manuscript.

331 All authors had full access to all of the data (including statistical reports and
332 tables) in the study and can take responsibility for the integrity of the data and
333 the accuracy of the data analysis.

334 Transparency declaration.

335 The lead author affirms that this manuscript is an honest, accurate, and
336 transparent account of the study being reported; that no important aspects of
337 the study have been omitted; and that any discrepancies from the study as
338 planned (and, if relevant, registered) have been explained.

339 Data sharing.

340 No additional data available.

341 **Sources of funding for the study.**

342 This research received funding from Conselho Nacional de Desenvolvimento
343 Científico e Tecnológico, Edital MCTI/CNPq/Universal 14/2014.

344 **Exclusive licence.**

345 “The Corresponding Author has the right to grant on behalf of all authors and
346 does grant on behalf of all authors, a worldwide licence to the Publishers and its
347 licensees in perpetuity, in all forms, formats and media (whether known now or
348 created in the future), to i) publish, reproduce, distribute, display and store the
349 Contribution, ii) translate the Contribution into other languages, create
350 adaptations, reprints, include within collections and create summaries, extracts
351 and/or, abstracts of the Contribution, iii) create any other derivative work(s)
352 based on the Contribution, iv) to exploit all subsidiary rights in the Contribution,
353 v) the inclusion of electronic links from the Contribution to third party material
354 where-ever it may be located; and, vi) licence any third party to do any or all of
355 the above.”

356 **Ethics approval.**

357 No protocol approval was necessary because data were obtained from
358 secondary sources.

359

Figure 1. The dietary content in added sugars regressed on the dietary contribution of ultra-processed foods evaluated by restricted cubic splines. US population aged 1 + years (NHANES 2009-2010) (N=9,317)

Legend: The values shown on the x-axis correspond to the 5th, 27.5th, 50th, 72.5th, and 95th percentiles for percentage of total energy from ultra-processed foods (knots). Coefficient for linear term=0.21 95% CI: 0.18 to 0.24. There was little evidence of nonlinearity in the restricted cubic spline model (Wald test for linear term $p<0.0001$; Wald test for all non-linear terms $p=0.19$).

REFERENCES

1. Guideline: Sugars intake for adults and children. Geneva: World Health Organization; 2015.

2. NHS Health Choices. Eight tips for healthy eating. <<http://www.nhs.uk/Livewell/Goodfood/Pages/eight-tips-healthy-eating.aspx>> (accessed August 2015).

3. Canadian Heart and Stroke Foundation Position Statement, August 2014. http://www.heartandstroke.com/site/c.iklQLcMWJtE/b.9201361/k.47CB/Sugar_heart_disease_and_stroke.htm (accessed August 2015)

4. Johnson RK, Appel LJ, Brands M, et al. on behalf of American Heart Association Nutrition Committee of the Council on Nutrition, Physical Activity, and Metabolism and the Council on Epidemiology and Prevention. Dietary sugars intake and cardiovascular health: a scientific statement from the American Heart Association. *Circulation* 2009;120(11):1011-1020. doi: 10.1161/CIRCULATIONAHA.109.192627

5. Scientific Report of the 2015 Dietary Guidelines Advisory Committee.

6. U.S. Department of Agriculture and U.S. Department of Health and Human Services. Dietary Guidelines for Americans, 2010. 7th Edition, Washington, DC: U.S. Government Printing Office, December 2010.

7. Marriott BP, Olsho L, Hadden L, et al. Intake of added sugars and selected nutrients in the United States, National Health and Nutrition Examination Survey (NHANES) 2003–2006. *Crit Rev Food Sci Nutr* 2010;50(3):228–58. doi: 10.1080/10408391003626223.

8. Bowman SA. Diets of individuals based on energy intakes from added sugars. *Family Economics and Nutrition Review* 1999;12(2):31–8.

9. Murphy S, Johnson R. The scientific basis of recent US guidance on sugars intake. *Am. J. Clin. Nutr.* 2003;78:827S-833S.

10. Welsh JA, Sharma AJ, Grellinger L, et al. Consumption of added sugars is decreasing in the United States. *Am J Clin Nutr* 2011;94:726–34. doi: 10.3945/ajcn.111.018366.

11. Yang Q, Zhang Z, Gregg EW, et al. Added Sugar Intake and Cardiovascular Diseases Mortality Among US Adults. *JAMA Intern Med.* 2014 Apr;174(4):516-24. doi:10.1001/jamainternmed.2013.13563.

12. United States Department of Agriculture. Economic Research Service. Sugar and Sweeteners Yearbook Tables. <<http://www.ers.usda.gov/data-products/sugar-and-sweeteners-yearbook-tables.aspx>> (accessed August 2015).

13. National Health and Nutrition Examination Survey <http://wwwn.cdc.gov/nchs/nhanes/search/nhanes09_10.aspx> (accessed August 2015).

14. Johnson CL, Paulose-Ram R, Ogden CL, et al. National Health and Nutrition Examination Survey: Analytic guidelines, 1999–2010. National Center for Health Statistics. *Vital Health Stat* 2 (161). 2013.

- http://www.cdc.gov/nchs/data/series/sr_02/sr02_161.pdf (accessed August 2015)
15. NHANES. MEC In-Person Dietary Interviewers Procedures Manual. January 2009.
<http://www.cdc.gov/nchs/data/nhanes/nhanes_09_10/DietaryInterviewers_Inperson.pdf> (accessed August 2015)
16. NHANES. Phone Follow-Up Dietary Interviewer Procedures Manual. September 2009.
<http://www.cdc.gov/nchs/data/nhanes/nhanes_09_10/Dietary_PFU_09.pdf> (accessed August 2015)
17. Moshfegh AJ, Rhodes DG, Baer DJ, et al. The USDA Automated Multiple-Pass Method reduces bias in the collection of energy intakes. *Am J Clin Nutr* 2008;88:324-332.
18. Blanton CA, Moshfegh AJ, Baer DJ, et al. The USDA Automated Multiple-Pass Method accurately estimates group total energy and nutrient intake. *J Nutr* 2006 Oct;136(10):2594-9.
19. Rumpler WV, Kramer M, Rhodes DG, et al. Identifying sources of reporting error using measured food intake. *Eur J Clin Nutr* 2008;62:544-52.
20. Automated Multiple-Pass Method. United States Department of Agriculture. Agriculture Research Service.
<http://www.ars.usda.gov/ba/bhnrc/fsrg> (accessed August 2015).
21. National Health and Nutrition Examination Survey. NHANES Response Rates and Population Totals. Response Rates.
<http://www.cdc.gov/nchs/nhanes/response_rates_CPS.htm> (accessed August 2015).
22. National Health and Nutrition Examination Survey. 2009 - 2010 Data Documentation, Codebook, and Frequencies.
http://www.cdc.gov/nchs/nhanes/nhanes2009-2010/DR1IFF_F.htm (accessed August 2015).
23. Monteiro CA, Levy RB, Claro RM, et al. A new classification of foods based on the extent and purpose of their processing. *Cad. Saúde Pública*, Nov 2010, vol.26, no.11, p.2039-2049.
24. Moubarac JC, Parra DC, Cannon G, et al. Food Classification Systems Based on Food Processing: significance and implications for policies and actions: a systematic literature review and assessment. *Curr Obes Rep* 2014;3:256-272.
25. Monteiro CA, Cannon G, Levy RB, et al. Ultra-processing and a new classification of foods. In: Neff R (ed) *Introduction to U.S. Food System. Public Health, Environment, and Equity*. San Francisco: Jossey Bass A Wiley Brand; 2015.
26. Monteiro CA. Nutrition and health. The issue is not food, nor nutrients, so much as processing. *Public Health Nutr*. 2009;12(5):729-731.

27. Ludwig DS. Technology, diet, and the burden of chronic disease. *JAMA* 2011;305:1352-1353.

28. Monteiro CA, Cannon G, Levy RB, et al. The Food System. Processing. The big issue for disease, good health, well-being. *World Nutrition* 2012; 3:527–569. www.wphna.org (accessed August 2015).

29. Moodie R, Stuckler D, Monteiro C, et al. Profits and pandemics: prevention of harmful effects of tobacco, alcohol, and ultra-processed food and drink industries. *Lancet*. 2013 Feb 23;381(9867):670-9. doi: 10.1016/S0140-6736(12)62089-3.

30. Ahuja JKA, Montville JB, Omolewa-Tomobi G, Heendeniya KY, Martin CL, Steinfeldt LC, Anand J, Adler ME, LaComb RP, and Moshfegh AJ. 2012. USDA Food and Nutrient Database for Dietary Studies, 5.0. U.S. Department of Agriculture, Agricultural Research Service, Food Surveys Research Group, Beltsville, MD.

31. U.S. Department of Agriculture, Agricultural Research Service. 2011. USDA National Nutrient Database for Standard Reference, Release 24. Nutrient Data Laboratory Home Page. <<http://www.ars.usda.gov/ba/bhnrc/ndl>> (accessed August 2015).

32. Bowman SA, Clemens JC, Thoeig RC, Friday JE, Shimizu M, and Moshfegh AJ. 2013. Food Patterns Equivalents Database 2009-10: Methodology and User Guide [Online]. Food Surveys Research Group, Beltsville Human Nutrition Research Center, Agricultural Research Service, U.S. Department of Agriculture, Beltsville, Maryland. <<http://www.ars.usda.gov/ba/bhnrc/fsrg>> (accessed June 2013).

33. Louzada MLC, Martins APB, Canella DC, et al. Ultra-processed foods and dietary quality in Brazil. *Rev. Saúde Pública São Paulo* 2015;vol.49; Epub July 10, 2015. doi:10.1590/S0034-8910.2015049006132

34. Dietary Guidelines for the Brazilian Population 2014. 2nd Edition. Ministry of Health of Brazil.


35. Eicher-Miller HA, Fulgoni III VL, Keast DR. Contributions of Processed Foods to Dietary Intake in the US from 2003–2008: A Report of the Food and Nutrition Science Solutions Joint Task Force of the Academy of Nutrition and Dietetics, American Society for Nutrition, Institute of Food Technologists, and International Food Information Council. *J. Nutr.* 2012;142:2065S–2072S. doi: 10.3945/jn.112.164442.

36. International Food Information Council Foundation. Understanding our food communications tool kit, 2010. <<http://www.foodinsight.org/For-Professionals/Understanding-Our-Food/tabid/1398/Default.aspx>> (accessed August 2015)

37. Poti JM, Mendez MA, Ng SW, et al. Is the degree of food processing and convenience linked with the nutritional quality of foods purchased by US households? *Am J Clin Nutr.* 2015 Jun;101(6):1251-62. doi:10.3945/ajcn.114.100925.

38. Moubarac J-C, Martins APB, Claro RM, et al. Consumption of ultra-processed foods and likely impact on human health. Evidence from Canada. *Public Health Nutr.* 2013 Dec;16(12):2240-8. doi: 10.1017/S1368980012005009.
39. Monteiro CA, Levy RB, Claro RM, et al. Increasing consumption of ultra-processed foods and likely impact on human health: evidence from Brazil. *Public Health Nutr.* 2011 Jan;14(1):5-13. doi:10.1017/S1368980010003241.
40. Crovetto M M, Uauy R, Martins AP, et al. Household availability of ready-to-consume food and drink products in Chile: impact on nutritional quality of the diet. *Rev Med Chil.* 2014 Jul; 142(7):850-8. doi:10.4067/S0034-98872014000700005.
41. Monteiro CA, Moubarac JC, Cannon G, et al. Ultra-processed products are becoming dominant in the global food system. *Obes Rev.* 2013 Nov;14 Suppl2:21-8. doi:10.1111/obr.12107.
42. Canella DS, Levy RB, Martins APB, et al. Ultra-processed food products and obesity in Brazilian households (2008-2009). *PLoS One.* 2014 Mar 25;9(3):e92752. doi:10.1371/journal.pone.0092752. eCollection 2014.
43. Bingham S, Luben R, Welch A, et al. Epidemiologic assessment of sugars consumption using biomarkers: comparisons of obese and nonobese individuals in the European Prospective Investigation of Cancer Norfolk. *Cancer Epidemiol Biomarkers Prev* 2007; 16(8):1651-4.
44. Lafay L, Mennen L, Basdevant A, et al. Does energy intake underreporting involve all kinds of food or only specific food items? Results from the Fleurbaix Laventie Ville Sante (FLVS) study. *Int J Obes Relat Metab Disord.* 2000; 24:1500-6.
45. Pryer JA, Vrijheid M, Nichols R, et al. Who are the "low energy reporters" in the dietary and nutritional survey of British adults? *Int J Epidemiol.* 1997; 26: 146-54.

Figure 1. The dietary content in added sugars regressed on the dietary contribution of ultra-processed foods evaluated by restricted cubic splines. US population aged 1 + years (NHANES 2009-2010) (N=9,317)



Online Supplementary Material

Food classification according to processing

Food items were initially classified into four groups shown in Table 1. This was accomplished by taking into account, the following three variables from the NHANES recall databases: "Main Food Description", "Additional Food Description" and "SR Code Description". Thereafter, the food item classification was modified, if necessary, taking two variables into account: "Combination Food Type" and "Source of food". Thus, most "Frozen meals" or "Lunchables" or food items consumed in "Restaurant fast food/pizza" or acquired at a "Vending machine", were classified as ultra-processed foods.

As explained in the Subjects and Methods section, when Food Codes were judged to be a handmade recipe, the classification was applied to the underlying ingredients (SR Codes), to enable a more precise food item classification (1).

It must be noted, however, that SR Codes and their proportions are not necessarily the ingredients and proportions consumed by the participant. One of the reasons is that links between FNDDS 5.0 and SR24 were developed to estimate the nutrient content of a Food Code and not the ingredient intake (2). Furthermore, when assigning SR Codes to a Food Code the individual-specific variable "Modification Code" ("adjustments to predefined recipe ingredients that reflect more closely the food as described by the respondent" (2)) was not taken into account, as manual changes would have had been necessary to do so.

Absence of data or discrepancies regarding degree of processing were solved opting for the lesser degree of processing (conservative criterion), which could have led to a slight underestimation of ultra-processed food consumption.

Online Supplementary Material

We classified homemade recipes with unknown ingredients based on expected principal ingredients, which could slightly underestimate ultra-processed food consumption.

Regarding bread, the classification distinguishes between handmade bread (either homemade or made in restaurants or artisanal bakeries), and industrial bread (made in industrial bakeries or factories), either processed (when made only of ingredients used in the making of handmade breads -flour, yeast, water, salt, and, sometimes, walnuts, dried fruits and other whole foods-) or ultra-processed (when adding substances not commonly used in the making of handmade breads -such as hydrogenated fat, sugars, starches, and additives). In our study, because of the large amount of industrial breads with unknown ingredients (approximately 3.7% of all industrial bread had fully known ingredients) and the very low consumption of processed breads when ingredients were reported (approximately 2.3% of industrial breads were processed), we ended up classifying all industrial bread as ultra-processed foods. This could slightly overestimate ultra-processed food consumption.

Assessing energy and added sugar contents

For some handmade recipes, the sum of the “calorie intake per SR Code” (calculated by us) of all underlying SR Codes did not add up exactly to the “calorie intake per Food Code” (provided by NHANES). In these cases, the “final calorie intake per SR code” was calculated as follows:

$$\text{Final calorie intake per SR code} = \text{NHANES Calorie intake per Food Code} * \left(\frac{\text{Calculated Calorie intake per SR code}}{\sum_{n=1}^{\infty} \text{Calculated Calorie intake per SR Code}} \right)$$

The same was done for added sugars:

Online Supplementary Material

$$\text{Final added sugars intake per SR code} = \text{Added sugars intake per Food Code} * \left(\frac{\text{Added sugars intake per SR code}}{\sum_{n=1}^{\infty} \text{Added sugars intake per SR code}} \right)$$

where n = each of the Food Code underlying SR Codes

For peer review only

Online Supplementary Material

Supplementary Table 1. NOVA food classification based on the extent and purpose of industrial processing (adapted from 3,4)

Food groups and definition	Examples
<p>1 Unprocessed or minimally processed foods</p> <p>Natural foods are those obtained directly from plants or animals (such as green leaves and fruits, or eggs and milk) and purchased for consumption without having undergone any alteration following their removal from nature. Minimally processed foods are natural foods that have been submitted to cleaning, removal of inedible or unwanted parts, fractioning, grinding, drying, fermentation, pasteurisation, cooling, freezing, or other processes which do not add substances to the original food. Purpose of minimum processes is to preserve foods and make it possible to store them and, sometimes, also to decrease stages of food preparation (cleaning and removing inedible parts) or facilitate their digestion, or render them more palatable (grinding or fermentation).</p>	<p>Natural, packaged, cut, chilled or frozen vegetables, fruits, potatoes, cassava, and other roots and tubers; bulk or packaged white, parboiled and wholegrain rice; whole or separated corn; grains of wheat and other cereals that are dried, polished, or ground as grits or flour; dried or fresh pasta made from wheat flour and water; all types of beans; lentils, chickpeas, and other legumes; dried fruits, fruit juices fresh or pasteurized without added sugar or other substances; nuts, peanuts, and other oilseeds without salt or sugar; fresh and dried mushrooms and other fungi; fresh and dried herbs and spices; fresh, frozen, dried beef, pork, poultry and other meat and fish; pasteurized, 'long-life' and powdered milk; fresh and dried eggs, yoghurt without sugar; and tea, herbal infusions, coffee, and tap, spring and mineral water.</p>
<p>2 Processed culinary ingredients</p> <p>These are substances extracted from natural foods or from nature itself by processes such as pressing, grinding, crushing, pulverising, and refining. Purpose of processing here is to obtain ingredients used in homes and restaurants kitchens to season and cook natural or minimally processed foods and to create with them varied and enjoyable dishes such as soups and broths, salads, rice and beans dishes, grilled or roasted vegetables and meat, and homemade breads, pies, cakes, and desserts.</p>	<p>Plant oils; coconut and animal fats (including butter and lard); table sugar, maple syrup (100%), molasses and honey; and table salt.</p>
<p>3 Processed foods</p> <p>These are relatively simple products manufactured essentially with the addition of salt or sugar or other substance of common culinary use, such as oil or vinegar, to natural or minimally processed foods. Purpose here is to prolong duration of foods and modify their palatability.</p> <p>If alcoholic beverages should be classified, drinks produced by the fermentation of group 1 food items such as wine, beer and cider will be classified in this group.</p>	<p>Canned and bottled vegetables, legumes or fruits; salted nuts or seeds; salted, smoked or cured meat or fish; canned sardine and tuna; cheeses, and breads made of wheat flour, yeast, water, and salt.</p>
<p>4 Ultra-processed foods</p> <p>These are food and drink products whose manufacture involves several stages and various processing techniques and ingredients, many of which are used exclusively by industry. Purpose of processing here is to create durable, accessible, convenient, and highly palatable, ready-to-drink, ready-to-eat, or ready-to-heat products typically consumed as snacks or desserts or as fast meals which replace dishes prepared from scratch.</p>	<p>Confectionery, soft drinks, sweetened juices and dairy drinks, powders for juices, sausages, chicken and fish nuggets or sticks and other pre-prepared frozen dishes, dried products such as cake mix, powdered soup, instant noodles, ready-seasonings, and an infinity of new products including packaged snacks, morning cereals, cereal bars, and 'energy' drinks. Sugar substitutes, sweeteners and all syrups (excluding 100% maple syrup). Breads and baked goods become ultra-processed products when, in addition to wheat flour,</p>

Online Supplementary Material

If alcoholic beverages should be classified, drinks produced by fermentation of group 1 food items followed by distillation and eventual addition of sugars or other substances, such as rum, whiskey, vodka, gin, and liqueurs, will be classified in this group.

yeast, water, and salt, their ingredients include substances such as hydrogenated vegetable fat, sugar, starch, whey, emulsifiers, and other additives.

For peer review only

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49

Online Supplementary Material

Supplementary Table 2. Percentage of participants with more than 10% of total energy intake from added sugars, by demographic subgroups, according to quintiles of the dietary contribution of ultra-processed foods. US population aged 1 + years (NHANES 2009-2010)

		Quintiles of the dietary contribution of ultra-processed foods (% of total energy intake)				
		1st (n=1955)	2nd (n=1901)	3rd (n=1774)	4th (n=1784)	5th (n=1903)
Gender	Men (n=4,634)	25.8	49.9	67.2	77.3	86.0*
	Women (n=4,683)	32.1	50.2	63.4	78.0	85.7*
Age (years)	1 to 5 (n=1,136)	23.8	47.6	60.1	71.5	90.7*
	6 to 11 (n=1,154)	39.5	59.6	71.6	84.5	91.0*
	12 to 19 (n=1,265)	43.1	69.1	73.4	78.0	89.0*
	20 to 39 (n=1,928)	26.7	52.7	71.5	76.7	86.9*
	40 to 59 (n=1,935)	32.1	46.6	59.5	79.6	81.7*
	60 and over (n=1,899)	24.6	44.4	60.4	72.6	74.7*
Race/ethnicity	Mexican American (n=2,064)	32.9	55.9	70.4	78.7	85.0*
	Other Hispanic (n=988)	37.7	55.4	72.4	80.9	90.6*
	Non-Hispanic White (n=3,984)	24.4	47.0	62.1	77.4	84.8*
	Non-Hispanic Black (n=1,726)	34.8	66.6	74.1	81.6	88.9*
	Other Race (including Multi-Racial) (n=555)	35.1	35.8	68.1	65.0	88.8*

Online Supplementary Material

Income to poverty*	0.00–1.30 (n=3,322)	33.4	59.8	73.6	83.9	86.9*
	>1.30–3.50 (n=3,062)	30.8	49.0	69.5	78.3	88.0*
	>3.50 and above (n=2,100)	22.6	45.0	54.5	73.3	81.6*

*Significant linear trend across quintiles ($P \leq 0.001$), both in unadjusted and Poisson models adjusted for sex, age group (1–5, 6–11, 12–19, 20–39, 40–59, 60 + years), race/ethnicity (Mexican-American, Other Hispanic, Non-Hispanic White, Non-Hispanic Black and Other Race - Including Multi-Racial-), and ratio of family income to poverty (SNAP 0.00–1.30, >1.30–3.50, and >3.50 and over).

Online Supplementary Material

References (for Online Supplementary Material).

1. Fitt E, Mak TN, Stephen AM, Prynne C, Roberts C, Swan G and Farron-Wilson M. Disaggregating composite food codes in the UK National Diet and Nutrition Survey food composition databank. *European Journal of Clinical Nutrition* (2010) 64, S32–S36.
2. Ahuja JKA, Montville JB, Omolewa-Tomobi G, Heendeniya KY, Martin CL, Steinfeldt LC, Anand J, Adler ME, LaComb RP, and Moshfegh AJ. 2012. USDA Food and Nutrient Database for Dietary Studies, 5.0. U.S. Department of Agriculture, Agricultural Research Service, Food Surveys Research Group, Beltsville, MD.
3. Moubarac JC, Parra DC, Cannon G, Monteiro CA. Food Classification Systems Based on Food Processing: significance and implications for policies and actions: a systematic literature review and assessment. *Curr Obes Rep* 2014; 3: 256-272.
4. Monteiro CA, Cannon G, Levy RB, Claro RM, Moubarac J-C. Ultra-processing and a new classification of foods. In: Neff R (ed) *Introduction to U.S. Food System. Public Health, Environment, and Equity*. San Francisco: Jossey Bass A Wiley Brand; 2015.

STROBE Statement—checklist of items that should be included in reports of observational studies

	Item No	Recommendation
Title and abstract	1	(a) Indicate the study's design with a commonly used term in the title or the abstract (Cross-sectional study; p.2) (b) Provide in the abstract an informative and balanced summary of what was done and what was found (p.2-3)
Introduction		
Background/rationale	2	Explain the scientific background and rationale for the investigation being reported (p.4-5)
Objectives	3	State specific objectives, including any prespecified hypotheses (p.5)
Methods		
Study design	4	Present key elements of study design early in the paper (p.6)
Setting	5	Describe the setting, locations, and relevant dates, including periods of recruitment, exposure, follow-up, and data collection (p.6-7)
Participants	6	(a) Cohort study—Give the eligibility criteria, and the sources and methods of selection of participants. Describe methods of follow-up (nap) Case-control study—Give the eligibility criteria, and the sources and methods of case ascertainment and control selection. Give the rationale for the choice of cases and controls (nap) Cross-sectional study—Give the eligibility criteria, and the sources and methods of selection of participants (p.6-7) (b) Cohort study—For matched studies, give matching criteria and number of exposed and unexposed (nap) Case-control study—For matched studies, give matching criteria and the number of controls per case (nap)
Variables	7	Clearly define all outcomes, exposures, predictors, potential confounders, and effect modifiers. Give diagnostic criteria, if applicable (p.7-10)
Data sources/measurement	8*	For each variable of interest, give sources of data and details of methods of assessment (measurement). Describe comparability of assessment methods if there is more than one group (p.7-10)
Bias	9	Describe any efforts to address potential sources of bias (lines 60-61)
Study size	10	Explain how the study size was arrived at (lines 54-60; lines 134-136)
Quantitative variables	11	Explain how quantitative variables were handled in the analyses. If applicable, describe which groupings were chosen and why (p.8-10)
Statistical methods	12	(a) Describe all statistical methods, including those used to control for confounding (p.9-10) (b) Describe any methods used to examine subgroups and interactions (p.9-10) (c) Explain how missing data were addressed (lines 134-136) (d) Cohort study—If applicable, explain how loss to follow-up was addressed (nap) Case-control study—If applicable, explain how matching of cases and controls was addressed (nap) Cross-sectional study—If applicable, describe analytical methods taking account of sampling strategy (lines 137-142) (e) Describe any sensitivity analyses (nap)

Continued on next page

Results		
Participants	13*	(a) Report numbers of individuals at each stage of study—eg numbers potentially eligible, examined for eligibility, confirmed eligible, included in the study, completing follow-up, and analysed (lines 54-61, lines 134-136) (b) Give reasons for non-participation at each stage (lines 54-61, lines 134-136) (c) Consider use of a flow diagram (nap)
Descriptive data	14*	(a) Give characteristics of study participants (eg demographic, clinical, social) and information on exposures and potential confounders (lines 147-158; Table 1) (b) Indicate number of participants with missing data for each variable of interest (lines 134-136) (c) Cohort study—Summarise follow-up time (eg, average and total amount) (nap)
Outcome data	15*	Cohort study—Report numbers of outcome events or summary measures over time (nap) Case-control study—Report numbers in each exposure category, or summary measures of exposure (nap) Cross-sectional study—Report numbers of outcome events or summary measures (lines 180-193; Table 1)
Main results	16	(a) Give unadjusted estimates and, if applicable, confounder-adjusted estimates and their precision (eg, 95% confidence interval). Make clear which confounders were adjusted for and why they were included (lines 194-224; Figure 1; Table 2) (b) Report category boundaries when continuous variables were categorized (Table 2) (c) If relevant, consider translating estimates of relative risk into absolute risk for a meaningful time period (nap)
Other analyses	17	Report other analyses done—eg analyses of subgroups and interactions, and sensitivity analyses (lines 220-221; Supplementary Table 2)
Discussion		
Key results	18	Summarise key results with reference to study objectives (lines 233-247)
Limitations	19	Discuss limitations of the study, taking into account sources of potential bias or imprecision. Discuss both direction and magnitude of any potential bias (lines 296-308)
Interpretation	20	Give a cautious overall interpretation of results considering objectives, limitations, multiplicity of analyses, results from similar studies, and other relevant evidence (lines 309-314)
Generalisability	21	Discuss the generalisability (external validity) of the study results (lines 265-288)
Other information		
Funding	22	Give the source of funding and the role of the funders for the present study and, if applicable, for the original study on which the present article is based (nap)

*Give information separately for cases and controls in case-control studies and, if applicable, for exposed and unexposed groups in cohort and cross-sectional studies.

Note: An Explanation and Elaboration article discusses each checklist item and gives methodological background and published examples of transparent reporting. The STROBE checklist is best used in conjunction with this article (freely available on the Web sites of PLoS Medicine at <http://www.plosmedicine.org/>, Annals of Internal Medicine at <http://www.annals.org/>, and Epidemiology at <http://www.epidem.com/>). Information on the STROBE Initiative is available at www.strobe-statement.org.

BMJ Open

Ultra-processed foods and added sugars in the US diet: evidence from a nationally representative cross-sectional study

Journal:	<i>BMJ Open</i>
Manuscript ID	bmjopen-2015-009892.R1
Article Type:	Research
Date Submitted by the Author:	29-Oct-2015
Complete List of Authors:	Martinez Steele, Euridice; University of Sao Paulo, Department of Nutrition, School of Public Health Baraldi, Larissa; University of Sao Paulo, Department of Nutrition, School of Public Health Louzada, Maria Laura; University of Sao Paulo, Department of Nutrition, School of Public Health; University of São Paulo, Center for Epidemiological Studies in Health and Nutrition Moubarac, Jean-Claude; University of São Paulo, Center for Epidemiological Studies in Health and Nutrition Mozaffarian, Dariush; Friedman School of Nutrition Science and Policy, Tufts University, Monteiro, Carlos; University of Sao Paulo, Department of Nutrition, School of Public Health; University of São Paulo, Center for Epidemiological Studies in Health and Nutrition
Primary Subject Heading:	Nutrition and metabolism
Secondary Subject Heading:	Public health
Keywords:	ultra-processed, added sugars, dietary intake, NHANES, US

SCHOLARONE™
Manuscripts

Ultra-processed foods and added sugars in the US diet: evidence from a nationally representative cross-sectional study

Names of authors:

Eurídice Martínez Steele (EMS; PHD student)^{ab},

Larissa Galastri Baraldi (LGB; PHD student)^{ab},

Maria Laura da Costa Louzada (ML; PHD student)^{ab},

Jean- Claude Moubarac (JCM, Researcher)^b,

Dariusz Mozaffarian (DM; Professor)^c,

Carlos Augusto Monteiro (CAM; Professor)^{ab}

^aDepartment of Nutrition, School of Public Health, University of São Paulo, São Paulo, Brazil

^bCenter for Epidemiological Studies in Health and Nutrition, University of São Paulo, São Paulo, Brazil

^cFriedman School of Nutrition Science and Policy, Tufts University, Boston, MA

Corresponding author:

Carlos Augusto Monteiro, Departamento de Nutrição, Faculdade de Saúde Pública, Universidade de São Paulo, Av. Dr. Arnaldo, 715, São Paulo 01246-907, Brazil. E-mail: carlosam@usp.br

Keywords: ultra-processed, added sugars, dietary intake, NHANES, US.

ABSTRACT

Objectives: To investigate the contribution of ultra-processed foods to the intake of added sugars in the US.

Ultra-processed foods were defined as industrial formulations which, besides salt, sugar, oils and fats, include substances not used in culinary preparations, in particular additives used to imitate sensorial qualities of minimally processed foods and their culinary preparations.

Design: Cross- sectional study.

Setting: *National Health and Nutrition Examination Survey 2009-2010.*

Participants: We evaluated 9,317 participants aged 1+ years with at least one 24-hour dietary recall.

Main outcome measures: Average dietary content of added sugars and proportion of individuals consuming more than 10% of total energy from added sugars.

Data analysis: Gaussian and Poisson regressions estimated the association between consumption of ultra-processed foods and intake of added sugars. All models incorporated survey sample weights and adjusted for age, sex, race/ethnicity, family income and educational attainment.

Results: Ultra-processed foods comprised 57.9% of energy intake, and contributed 89.7% of the energy intake from added sugars. The content of added sugars in ultra-processed foods (21.1% of calories) was 8-fold higher than in processed foods (2.4%) and 5-fold higher than in unprocessed or

minimally processed foods and processed culinary ingredients grouped together (3.7%). In both unadjusted and adjusted models, each increase of 5 percentage points in proportional energy intake from ultra-processed foods increased the proportional energy intake from added sugars by 1 percentage point. Consumption of added sugars increased linearly across quintiles of ultra-processed food consumption: from 7.5% of total energy in the lowest quintile to 19.5% in the highest. A total of 82.1% of Americans in the highest quintile exceeded the recommended limit of 10% energy from added sugars, compared with 26.4% in the lowest.

Conclusions: Decreasing the consumption of ultra-processed foods could be an effective way of reducing the excessive intake of added sugars in the US.

Strengths and limitations of this study:

- Use of a large, nationally representative sample of the US population, increasing generalizability.
- Use of data on added sugars rather than total sugars or sugar-sweetened beverages, which corresponds to guidelines relevant area of prioritization.
- Unlike most articles which have focused on specific food items such as soft drinks or fast food, our study evaluates the impact of a comprehensive group of products whose consumption is increasing exponentially in most countries.
- Dietary data obtained by 24-hour recalls is subjected to potential error and bias.

- Information indicative of food processing is not consistently determined for all food items in NHANES, which could lead to modest over or underestimation of the consumption of ultra-processed foods.

For peer review only

1 INTRODUCTION

2 Increasing policy attention has focused on added sugars, including by the World
3 Health Organization (WHO)(1), the United Kingdom National Health System(2),
4 the Canadian Heart and Stroke Foundation(3), the American Heart Association
5 (AHA)(4), and the US Dietary Guidelines Advisory Committee (USDGAC)(5).

6 These reports concluded that a high intake of added sugars increases risk for
7 weight gain(1,4,5), excess body weight(5) and obesity(3,5); type 2 diabetes
8 mellitus(3,5); higher serum triglycerides(5) and high blood cholesterol(3); higher
9 blood pressure(5) and hypertension(5); stroke(3,5); coronary heart disease(3,5);
10 cancer(3); and dental caries(1,3,5). Moreover, foods higher in added sugars
11 are often a source of empty calories with minimum essential nutrients or dietary
12 fiber(6-8), which displace more nutrient-dense foods(9) and lead, in turn, to
13 simultaneously overfed and undernourished individuals.

14 All reports recommended limiting intake of added sugars(1,3-5). In the US, the
15 USDGAC recommended limiting added sugars to no more than 10% of total
16 calories. This is a challenge, as recent consumption of added sugars in the US
17 amounted to almost 15% of total calories in 2005-2010(10,11).

18 To design and implement effective measures to reduce added sugars, their
19 dietary sources must be clearly identified. Added sugars can be consumed
20 either as ingredients of dishes or drinks prepared from scratch by consumers or
21 cook, or as ingredients of food products manufactured by the food
22 industry. According to market disappearance data from 2014, more than three
23 quarters of the sugar and high fructose corn syrup available for human
24 consumption in the US were used by the food industry(12). This suggests food

1
2
3 25 products manufactured by the industry could have an important role in the
4
5 26 excess added sugars consumption in the US. However, to assess this role, it is
6
7 27 essential to consider the contribution of manufactured food products to both
8
9 28 total energy intake and the energy intake from added sugars, and, more
10
11 29 relevantly, to quantify the relationship between their consumption and the total
12
13 30 dietary content of added sugars. To address these questions, we performed an
14
15 31 investigation utilizing 2009-2010 *National Health and Nutrition Examination*
16
17 32 *Survey (NHANES)*.
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

33 **SUBJECTS AND METHODS**

34 **Data source, population and sampling**

35 We utilized nationally representative data from the 2009-2010 *National Health*
36 *and Nutrition Examination Survey (NHANES)*, specifically the dietary
37 component *What we eat in America (WWEIA)*(13).

38 NHANES is a continuous, nationally representative, cross-sectional survey of
39 the non-institutionalized, civilian US residents(14). NHANES sample was
40 obtained by using a complex, stratified, multi-stage probability cluster sampling
41 design, based on the selection of counties, blocks, households, and the number
42 of people within households(14). In order to improve the estimate precision and
43 reliability, NHANES 2009-2010 oversampled the following subgroups: Hispanic,
44 Non-Hispanic black, Non-Hispanic white and Other persons at or below 130%
45 of the federal poverty level and Non-Hispanic white and Other persons aged 80
46 + years(14).

47 The survey included an interview conducted in the home and a subsequent
48 health examination performed at a mobile examination center (MEC). All
49 NHANES examinees were eligible for two 24-hour dietary recall interviews. The
50 first dietary recall interview was collected in-person in the MEC(15) while the
51 second was collected by telephone 3 to 10 days later but never on the same
52 day of the week as the MEC interview(16). Dietary interviews were conducted
53 by trained interviewers using the validated(17-19) *US Department of Agriculture*
54 *Automated Multiple-Pass Method (AMPM)*(20). For children under 9 years of
55 age, the interview was conducted with a proxy; for children between 6 and 8
56 years of age, in the presence of the child. Children 9 to 11 years old provided

their own data assisted by an adult household member (assistant). The preferred proxy/assistant was the most knowledgeable person about the child's consumption the day before the interview. If the child had more than one caregiver, several individuals could contribute to the intake data(15; 16).

Among the 13,272 people screened in NHANES 2009-2010, 10,537 (79.4%) participated in the household interview and 10253 (77.3%) also participated in the MEC health examination(21). Of these, 9,754 individuals provided one day of complete dietary intakes and 8,406 provided two days(22).

We evaluated 9,317 survey participants aged 1 year and above who had one day 24-hour dietary recall data and had not been breast-fed on either of the two days. These individuals had similar socio-demographic characteristics (gender, age, race/ ethnicity, family income and educational attainment) to the full sample of 10,109 participants interviewed.

Food classification according to processing

We classified all recorded food items (N=280,132 Food Codes for both recall days) according to NOVA, a food classification based on the extent and purpose of industrial food processing(23-25). This classification includes 4 groups: "unprocessed or minimally processed foods" (such as fresh, dry or frozen fruits or vegetables, grains, legumes, meat, fish and milk); "processed culinary ingredients" (including table sugar, oils, fats, salt, and other substances extracted from foods or from nature, and used in kitchens to make culinary preparations), "processed foods" (foods manufactured with the addition of salt or sugar or other substances of culinary use to unprocessed or minimally processed foods, such as simple cheese, bread and canned food), and "ultra-

processed foods” (formulations of several ingredients which, besides salt, sugar, oils and fats, include food substances not used in culinary preparations. In particular flavors, colors, sweeteners, emulsifiers and other additives used to imitate sensorial qualities of unprocessed or minimally processed foods and their culinary preparations or to disguise undesirable qualities of the final product). A detailed definition of each food group and examples of food items classified in each group are shown in **Supplementary Table 1**. The rationale underlying the classification is described elsewhere(26-29).

For all food items (Food Codes) judged to be a handmade recipe, the classification was applied to the underlying ingredients (Standard Reference Codes -SR Codes-) obtained from the USDA Food and Nutrient Database for Dietary Studies (FNDDS) 5.0(30). Refer to Online Supplementary Material (OSM) for further details.

Assessing energy and added sugar contents

For this study, we used Food Code energy values as provided by NHANES.

For handmade recipes, we calculated the underlying ingredient (SR Code) energy values using variables from both *FNDDS 5.0*(30) and *USDA National Nutrient Database for Standard Reference, Release 24 (SR24)*(31). Refer to OSM for further details.

Data on added sugars per Food Code and per SR Code were obtained by merging the Food Patterns Equivalents Database (FPED) 2009-2010 and the Food Patterns Equivalents Ingredients Database (FPID) 2009-2010(32). Added sugars are defined in these databases as “sugars that are added to foods as an ingredient during preparation, processing, or at the table. Added sugars do not

1
2
3 105 include naturally occurring sugars (e.g., lactose in milk, fructose in fruits).
4
5 106 Examples of added sugars include brown sugar, cane sugar, confectioners'
6
7 107 sugar, granulated sugar, dextrose, white sugar, corn syrup and corn syrup
8
9 108 solids, molasses, honey, and all types of syrups such as maple syrup, table
10
11 109 syrups, and pancake syrup”(32). These two databases express the content of
12
13 110 added sugars in teaspoons per 100 g. Teaspoons were converted into grams
14
15 111 using the factor 4.2 g/teaspoon and into kcal using the factor 3.87 kcal/g.
16
17
18

19 112 **Data Analysis**

20
21
22 113 We utilized all available day 1 dietary data for each participant. Food items
23
24 114 were sorted into mutually exclusive food subgroups within Unprocessed or
25
26 115 minimally processed foods (n= 11), Processed culinary ingredients (n=4),
27
28 116 Processed foods (n=4) and Ultra-processed foods (n=18), as shown in Table 1.
29
30 117 First, we evaluated the contributions of each of the NOVA food groups and
31
32 118 subgroups to total energy and to the energy from added sugars. Next, we
33
34 119 calculated the average content of added sugars in the overall US diet and in
35
36 120 fractions of this diet composed by each of the NOVA food groups and
37
38 121 subgroups. We also calculated the dietary content of added sugars in the group
39
40 122 of unprocessed or minimally processed foods combined with the group of
41
42 123 processed culinary ingredients, as foods belonging to these two groups are
43
44 124 usually combined together in culinary preparations and therefore consumed
45
46 125 together.
47
48
49

50
51
52 126 We used Gaussian regression to estimate the association between the dietary
53
54 127 contribution of ultra-processed foods and the dietary content of added sugars,
55
56 128 each expressed as proportions of total energy. This association was also
57
58
59
60

139 explored after adjusting for the proportion of added sugars in non-ultra-
130 processed energy intake. Dietary contribution of ultra-processed foods was
131 transformed using restricted cubic spline functions to allow for nonlinearity.

132 The average content of added sugars in the overall diet was compared across
133 quintiles of the dietary contribution of ultra-processed foods. Poisson
134 regression was used to assess whether the percentage of diets with more than
135 10% or 20% of total energy from added sugars increased across quintiles. This
136 increase was also evaluated across demographic subgroups in stratified
137 analysis. Tests of linear trend were performed in order to evaluate the effect of
138 quintiles as a single continuous variable.

139 All regression models were adjusted for age (1-5 years, 6-11 years, 12-19
140 years, 20–39 years, 40–59 years, 60 + years), sex, race/ethnicity (Mexican-
141 American, Other Hispanic, Non-Hispanic White, Non-Hispanic Black, Other
142 Race including Multi-Racial), ratio of family income to poverty (categorized
143 based on Supplemental Nutrition Assistance Program (SNAP) eligibility as
144 0.00–1.30, >1.30–3.50, and >3.50 and above)(14) and educational attainment
145 of respondents, for participants aged 20 + years, and of household reference
146 person otherwise (<12, 12 years and >12 years). As 908 participants had
147 missing values on family income and/or educational attainment, multivariable-
148 adjusted analysis included 8,409 individuals. Analysis which also adjusted for
149 the added sugar content of all non-ultra-processed foods grouped together
150 included 8,335 individuals.

151 NHANES survey sample weights were used in all analyses to account for
152 differential probabilities of selection for the individual domains, nonresponse to

1
2
3 153 survey instruments, and differences between the final sample and the total US
4
5 154 population. The Taylor series linearization variance approximation procedure
6
7 155 was used for variance estimation in all analysis in order to account for the
8
9 156 complex sample design and the sample weights(14).
10
11
12 157 To minimize chance findings from multiple comparisons, statistical hypotheses
13
14 158 were tested using a two-tailed $p < 0.001$ level of significance. Data were
15
16
17 159 analyzed using Stata statistical software package version 12.1.
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

160 **RESULTS**

161 **Distribution of total energy intake by food groups**

162 The average US daily energy intake in 2009-2010 was 2069.5 kcal, and nearly
163 3 in 5 calories (57.9%) came from ultra-processed foods (**Table 1**).

164 Unprocessed or minimally processed foods contributed 29.6% of total calories,
165 processed foods an additional 9.4%, and processed culinary ingredients the
166 remaining 2.9%. The most common ultra-processed foods in terms of energy
167 contribution were breads; soft drinks, fruit drinks, and milk-based drinks; cakes,
168 cookies, and pies; salty-snacks; frozen and shelf-stable plates; pizza; and
169 breakfast cereals. Meat, fruit, and milk provided the most calories among
170 unprocessed or minimally processed foods; ham and cheese, the most calories
171 among processed foods; and table sugar and plant oils, the most calories
172 among processed culinary ingredients.

173

Table 1. Distribution of the total energy intake and of the energy intake from added sugars according to food groups, and the mean content of added sugars of each food group. US population aged 1 + years (NHANES 2009-2010) (N=9,317)

Food groups	Mean energy intake		Mean energy intake from added sugars		Mean content of added sugars
	Absolute (kcal/day)	Relative (% of total energy intake)	Absolute (kcal/day)	Relative (% of total energy intake from added sugars)	% of energy from added sugars
Unprocessed or minimally processed foods	585.5	29.6	0.0	0.0	0.0
Meat (includes poultry)	165.3	7.9	0.0	0.0	0.0
Fruit ¹	97.5	5.2	0.0	0.0	0.0
Milk and plain yoghurt	96.4	5.1	0.0	0.0	0.0
Grains	53.3	2.8	0.0	0.0	0.0
Roots and tubers	32.2	1.6	0.0	0.0	0.0
Eggs	28.8	1.4	0.0	0.0	0.0
Pasta	28.4	1.4	0.0	0.0	0.0
Legumes	16.2	0.8	0.0	0.0	0.0
Fish and sea food	17.2	0.8	0.0	0.0	0.0
Vegetables	13.5	0.7	0.0	0.0	0.0
Other unprocessed or minimally processed foods ²	36.7	1.8	0.0	0.0	0.0
Processed culinary ingredients	64.3	2.9	24.4	8.7	38.8

Table sugar ³	24.7	1.1	24.4	8.7	98.5
Plant oils	27.5	1.3	0.0	0.0	0.0
Animal fats ⁴	11.2	0.5	0.0	0.0	0.0
Other processed culinary ingredients ⁵	0.9	0.04	0.0	0.0	0.0
Unprocessed or minimally processed foods + Processed culinary ingredients	649.8	32.6	24.4	8.7	3.7
Processed foods	209.7	9.4	2.5	1.6	2.4
Cheese	80.1	3.7	0.0	0.0	0.0
Ham and other salted, smoked or canned meat or fish	26.4	1.2	0.3	0.2	1.4
Vegetables and other plant foods preserved in brine	13.4	0.7	1.6	0.9	13.7
Other processed foods ⁶	89.8	3.8	0.6	0.5	1.2
Ultra-processed foods	1209.8	57.9	265.2	89.7	21.1
Breads ⁷	191.6	9.5	10.6	7.6	5.7
Cakes, cookies and pies	122.8	5.7	29.8	11.2	24.2
Salty-snacks	93.2	4.6	1.2	0.7	1.4
Frozen and shelf-stable plate meals	80.6	4.02	1.1	0.7	1.6
Soft drinks, carbonated	81.8	3.7	75.2	17.1	69.9
Pizza (ready-to-eat/heat)	81.8	3.5	2.4	1.4	2.9
Fruit drinks ⁸	69.2	3.3	55.7	13.9	67.5

Breakfast cereals	50.9	2.8	12.4	6.4	23.3
Sauces, dressings and gravies	49.8	2.4	4.4	2.8	10.0
Reconstituted meat or fish products	51.5	2.4	0.7	0.6	2.0
Sweet-snacks	50.9	2.4	19.4	7.1	38.9
Ice cream and ice pops	48.7	2.3	18.3	5.9	36.9
Milk-based drinks ⁹	34.6	1.8	10.8	4.6	34.1
Desserts ¹⁰	36.4	1.8	18.5	7.3	48.5
French fries and other potatoe products ¹¹	37.8	1.7	0.0	0.0	0.0
Sandwiches and hamburgers on bun (ready-to-eat/heat)	32.5	1.4	1.3	0.6	4.4
Instant and canned soups	14.3	0.8	0.1	0.1	0.7
Other ultra-processed foods ¹²	81.5	3.8	3.1	1.5	7.8
Total	2069.5	100.0	292.2	100.0	13.8

176 1Including freshly squeezed juices

177 2Including nuts and seeds (unsalted); yeast; dried fruits (without added sugars) and vegetables; non pre-sweetened, non-whitened, non-
178 flavored coffee and tea; coconut water and meat; homemade soup and sauces; flours; tapioca

179 3Including honey, molasses, maple syrup (100%)

180 4Including unsalted butter, lard and cream

181 5Including starches; coconut and milk cream; unsweetened baking chocolate, cocoa powder and gelatin powder; vinegar; baking powder and
182 baking soda

183 6Including salted or sugared nuts and seeds; peanut, sesame, cashew and almond butter or spread; beer and wine

- 1
- 2
- 3
- 4
- 5
- 61847Including all types of bread. Processed bread made of flour, water, salt, leavening agents and possibly walnuts, dried fruits and other whole
- 7185foods, were included under this group as well, because of the low consumption
- 8
- 91868Including fruit and fruit-flavored, non-carbonated and other sweetened drinks, including presweetened tea and coffee, energy drinks, sports
- 10187drinks with no milk added, nonalcoholic wine
- 11
- 121889Including flavored yogurt sweetened with sugar or with low-calorie sweetener, milk-shake, soymilk
- 13
- 1418910Including ready-to-eat and dry-mix desserts such as pudding; sweetened canned fruit and fruit sauce
- 15
- 1619011Including hash browns, potato puffs, stuffed potatoes, onion rings (ready-to-eat/heat)
- 17
- 1819112Including soy products such as meatless patties and fish sticks; babyfood and baby formula; dips, spreads, mustard and catsup; salted butter
- 19192and margarine; sugar substitutes, sweeteners and all syrups (excluding 100% maple syrup); distilled alcoholic drinks
- 20
- 21
- 22
- 23
- 24
- 25
- 26
- 27
- 28
- 29
- 30
- 31
- 32
- 33
- 34
- 35
- 36
- 37
- 38
- 39
- 40
- 41
- 42
- 43
- 44
- 45
- 46
- 47
- 48
- 49

193 **Distribution of energy intake from added sugars by food groups**

194 The average US daily intake of added sugars was 292.2 kcal (Table 1).
195 Notably, almost 90% of this (89.7%) came from ultra-processed foods. The
196 main sources of added sugars among ultra-processed foods were: soft drinks
197 (17.1% of US intake of added sugars), fruit drinks (13.9%), milk-based drinks
198 (4.6%); cakes, cookies, and pies (11.2%); breads (7.6%); desserts (7.3%);
199 sweet snacks (7.1%); breakfast cereals (6.4%); and ice creams and ice pops
200 (5.9%). In contrast, only 8.7% of the added sugars in the US diet came from
201 processed culinary ingredients (table sugar consumed as part of dishes or
202 drinks prepared from scratch by consumers or cook), and only 1.6% from
203 processed foods.

204 The average content of added sugars in ultra-processed foods (21.1% of
205 calories) was 8-fold higher than in processed foods (2.4%) and 5-fold higher
206 than in unprocessed or minimally processed foods and processed culinary
207 ingredients grouped together (3.7%) (Table 1).

208 **Association between consumption of ultra-processed foods and added** 209 **sugar intake**

210 In unadjusted restricted cubic splines Gaussian regression analysis, a strong
211 linear association was identified between the dietary contribution (percentage of
212 calories) of ultra-processed foods and the dietary content (percentage of
213 calories) in added sugars (coefficient for linear term=0.20, 95% CI: 0.17 to 0.23)
214 **(Figure 1).**

215 **Figure 1.**

216 There was little evidence of nonlinearity in the restricted cubic spline model
217 (Wald test for linear term $p<0.0001$; Wald test for all non-linear terms $p=0.27$).
218 The strength of the association remained fairly the same after adjusting for age,
219 sex, race/ethnicity, family income, educational attainment and proportion of
220 added sugars in non-ultra-processed energy intake (coefficient for linear
221 term=0.19, 95% CI: 0.17 to 0.22). Overall, each increase in 5 percentage points
222 of energy in consumption of ultra-processed foods was associated with 1 higher
223 percentage point of energy in the consumption of added sugars.

224 Across quintiles of energy-adjusted ultra-processed food consumption, the
225 intake of added sugars increased substantially and monotonically, from 7.5% of
226 total calories in the lowest quintile to 19.5% in the highest. Across the same
227 quintiles, the proportion of individuals consuming more than 10% of total energy
228 from added sugars (59.6% in the total population) increased from 26.4% to
229 82.1%, respectively. An even more pronounced increase was seen in the
230 proportion of individuals consuming more than 20% of their total energy from
231 added sugars: from 4.7% in the lowest quintile to 41.2% in the highest (**Table**
232 **2**). Similar increases were seen in stratified analysis by major demographic
233 subgroups (**Supplementary Table 2**). The magnitude and the statistical
234 significance of the association between the dietary contribution of ultra-
235 processed foods and the dietary content in added sugars did not change with
236 adjustment for sex, age, race/ethnicity, family income and educational
237 attainment.

Table 2. Indicators of the dietary content in added sugars according to the dietary contribution of ultra-processed foods. US population aged 1 + years (NHANES 2009-2010)

		Indicators						
Dietary contribution of ultra-processed foods (% of total energy intake)		% of total energy intake from added sugars	Participants with more than 10% of total energy intake from added sugars			Participants with more than 20% of total energy intake from added sugars		
Quintiles	Mean (range)	Mean	%	PR ¹	PRadj ²	%	PR ¹	PRadj ²
1st (n=1,937)	28.9 (0 to 40.1)	7.5	26.4	1	1	4.7	1	1
2nd (n=1,888)	47.3 (40.1 to 53.3)	11.1	50	1.9	1.9	10.5	2.2	2.2
3rd (n=1,814)	58.7 (53.3 to 64.1)	13.8	62.7	2.4	2.3	21.1	4.5	4.3
4th (n=1,779)	69.7 (64.1 to 75.7)	16.9	76.6	2.9	2.8	29.9	6.4	5.9
5th (n=1,899)	85.1 (75.7 to 100)	19.5*	82.1	3.1*	2.9*	41.2	8.8*	7.9*
Total (n=9,317)	57.9 (0 to 100.0)	13.8	59.6	—	—	21.5	—	—

*Significant linear trend across all quintiles ($p \leq 0.001$), both in unadjusted and models adjusted for sex, age group (1-5, 6-11, 12-19, 20-39, 40-59, 60 + years), race/ethnicity (Mexican-American, Other Hispanic, Non-Hispanic White, Non-Hispanic Black and Other Race - Including Multi-Racial-), ratio of family income to poverty (SNAP 0.00-1.30, >1.30-3.50, and >3.50 and over) and educational attainment (<12, 12 years and >12 years).

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49

244 1PR=Prevalence ratios estimated using Poisson regression (N=9,317)

245 2PRadj=Prevalence ratios adjusted for sex, age groups, race/ethnicity, ratio of family income to poverty and educational attainment,

246 as above (N=8,409)

For peer review only

DISCUSSION

In this analysis of nationally representative data, we confirmed the excessive consumption of added sugars in the US(10,11). We also provide new evidence that ultra-processed foods represent more than half of all calories in the US diet, and contribute nearly 90% of all added sugars. Added sugars represented 1 of every 5 calories in the average ultra-processed food (21.1%), far higher than the content of added sugars in processed foods (2.4%) and in unprocessed or minimally processed foods, and processed culinary ingredients grouped together (3.7%). A strong linear relationship was found between the dietary contribution of ultra-processed foods and the dietary content of added sugars. Moreover, the risk of exceeding the recommended upper limit of 10% energy from added sugars was far higher when ultra-processed food consumption was high, and risk differences were even more pronounced for exceeding a limit of 20% energy. Notably, only those Americans in the lowest quintile of ultra-processed food consumption met the recommended limit of <10% energy from added sugars. To our knowledge, this is the first study to assess the consumption of ultra-processed foods and establish its relationship with excessive added sugar intake in the US.

The high consumption of added sugars in the US is likely contributing to excess obesity, type 2 diabetes, dyslipidemia, hypertension and coronary heart disease(1,3-5). Consequently, most dietary guidelines now recommend limiting added sugar consumption. However, such guidelines are not always clear on how to put this recommendation into practice. Our study suggests that in the US, limiting the consumption of ultra-processed foods may be a highly effective way to decrease added sugars. A reduction in ultra-processed foods should

272 also increase the intake of more healthful, minimally processed foods such as
273 milk, fruits, and nuts, and freshly-prepared dishes based on whole grains and
274 vegetables, which would produce additional health benefits beyond the
275 reduction in added sugar. Consistent with this approach, in Brazil, where the
276 consumption of added sugars is as high as in the US(33), the new dietary
277 guidelines launched in 2014 emphasize the importance of not replacing
278 unprocessed or minimally processed foods and freshly prepared dishes by
279 ultra-processed foods(34).

280 Few studies have assessed the impact of levels of food processing on the
281 nutrient profile of the US diet. One analysis using data from NHANES 2003-
282 2008(35) used a food classification system(36) including “Mixtures of combined
283 Ingredients” and “Ready-to-eat”, which are mostly ultra-processed foods and
284 together, contributed to about half of total energy intake and three-quarters of
285 energy intake from added sugars. Another study evaluated household
286 barcoded purchasing data from 2000-2012 using a classification system guided
287 by the one used in our study(37). In 2012, the mean per capita purchase of
288 “highly processed foods”, a category similar to ultra-processed foods,
289 corresponded to 61.0% of all calories and had higher adjusted median total
290 sugar content than “less processed foods”. This report did not evaluate added
291 sugars nor the contribution of processed foods to sugar intake. It also did not
292 capture non-barcoded items such as unpackaged fresh fruit, vegetables and
293 meat, or highly processed foods such as ready-to-eat store-prepared items. An
294 investigation in Canada, using 2001 household purchasing data, found that
295 ultra-processed foods are high in free sugars and that only households in the
296 lowest quintile of ultra-processed food purchasing might have met the

recommended limit of <10% energy from free sugars (9.2%)(38). Being based on household purchasing data, these two prior studies and others based on the NOVA classification system(23, 39-42) could not evaluate fraction of wasted food nor purchases at restaurants, which represent a substantial proportion of US calories. Our findings build upon and considerably extend these prior reports by evaluating food processing and added sugar intake using contemporary, nationally representative dietary intake data in the US.

Our study has several strengths. We studied a large, nationally representative sample of the US population, increasing generalizability. Use of data on added sugars rather than total sugars or sugar-sweetened beverages, corresponds to the relevant area of prioritization of recent national and international guidelines. Our investigation was based on individual consumption data, rather than market disappearance or household purchasing data which cannot account for differences between amounts purchased and amounts actually consumed.

Potential limitations should be considered. As with most population measures, dietary data obtained by 24-hour recalls is imperfect. However, the standardized methods and approach of NHANES minimize potential error and bias, particularly for assessing population averages as focused upon in the present study. Previous studies suggest that people with obesity may underreport consumption of foods with caloric sweeteners(43) such as desserts and sweet baked goods(44, 45). If so, these biases may lead to an underestimation of the dietary contribution of ultra-processed foods and the overall intake of added sugars, but should have much less effect on the association between these. Although NHANES collects some information indicative of food processing (i.e. place of meals, product brands), these data

are not consistently determined for all food items, which could lead to modest over or underestimation of the consumption of ultra-processed foods.

In conclusion, we found that ultra-processed foods contribute almost 60% of calories and 90% of added sugars consumed in the US. Only Americans in the lowest quintile of ultra-processed food consumption met the recommended guidelines for intake of added sugars. Decreasing the consumption of ultra-processed foods could be an effective way of reducing the excessive intake of added sugars in the US.

330

331

Competing Interests.

"All authors have completed the ICMJE uniform disclosure form at www.icmje.org/coi_disclosure.pdf and declare: no support from any organisation for the submitted work; DM reports ad hoc honoraria or consulting from Bunge, Haas Avocado Board, Nutrition Impact, Amarin, Astra Zeneca, Boston Heart Diagnostics, GOED, and Life Sciences Research Organization; and scientific advisory boards, Unilever North America and Elysium Health; no other relationships or activities that could appear to have influenced the submitted work."

Contributorship statement.

CAM, EMS, DM designed research; EMS, LB, ML data management; EMS, J-CM, ML analyzed data; EMS, DM, CAM wrote paper; CAM, EMS had primary responsibility for final content. All authors read and approved the final manuscript.

All authors had full access to all of the data (including statistical reports and tables) in the study and can take responsibility for the integrity of the data and the accuracy of the data analysis.

Transparency declaration.

The lead author affirms that this manuscript is an honest, accurate, and transparent account of the study being reported; that no important aspects of the study have been omitted; and that any discrepancies from the study as planned (and, if relevant, registered) have been explained.

Data sharing.

355 No additional data available.

356 **Sources of funding for the study.**

357 This research received funding from Conselho Nacional de Desenvolvimento
358 Científico e Tecnológico, Edital MCTI/CNPq/Universal 14/2014.

359 **Exclusive licence.**

360 “The Corresponding Author has the right to grant on behalf of all authors and
361 does grant on behalf of all authors, a worldwide licence to the Publishers and its
362 licensees in perpetuity, in all forms, formats and media (whether known now or
363 created in the future), to i) publish, reproduce, distribute, display and store the
364 Contribution, ii) translate the Contribution into other languages, create
365 adaptations, reprints, include within collections and create summaries, extracts
366 and/or, abstracts of the Contribution, iii) create any other derivative work(s)
367 based on the Contribution, iv) to exploit all subsidiary rights in the Contribution,
368 v) the inclusion of electronic links from the Contribution to third party material
369 where-ever it may be located; and, vi) licence any third party to do any or all of
370 the above.”

371 **Ethics approval.**

372 No protocol approval was necessary because data were obtained from
373 secondary sources.

374

Figure 1. The dietary content in added sugars regressed on the dietary contribution of ultra-processed foods evaluated by restricted cubic splines. US population aged 1 + years (NHANES 2009-2010) (N=9,317)

Legend: The values shown on the x-axis correspond to the 5th, 27.5th, 50th, 72.5th, and 95th percentiles for percentage of total energy from ultra-processed foods (knots). Coefficient for linear term=0.20 95% CI: 0.17 to 0.23. There was little evidence of nonlinearity in the restricted cubic spline model (Wald test for linear term $p<0.0001$; Wald test for all non-linear terms $p=0.27$).

REFERENCES

1. Guideline: Sugars intake for adults and children. Geneva: World Health Organization; 2015.
2. NHS Health Choices. Eight tips for healthy eating. <<http://www.nhs.uk/Livewell/Goodfood/Pages/eight-tips-healthy-eating.aspx>> (accessed August 2015).
3. Canadian Heart and Stroke Foundation Position Statement, August 2014. http://www.heartandstroke.com/site/c.iklQLcMWJtE/b.9201361/k.47CB/Sugar_heart_disease_and_stroke.htm (accessed August 2015)
4. Johnson RK, Appel LJ, Brands M, et al. on behalf of American Heart Association Nutrition Committee of the Council on Nutrition, Physical Activity, and Metabolism and the Council on Epidemiology and Prevention. Dietary sugars intake and cardiovascular health: a scientific statement from the American Heart Association. *Circulation* 2009;120(11):1011-1020. doi: 10.1161/CIRCULATIONAHA.109.192627
5. Scientific Report of the 2015 Dietary Guidelines Advisory Committee.
6. U.S. Department of Agriculture and U.S. Department of Health and Human Services. Dietary Guidelines for Americans, 2010. 7th Edition, Washington, DC: U.S. Government Printing Office, December 2010.
7. Marriott BP, Olsho L, Hadden L, et al. Intake of added sugars and selected nutrients in the United States, National Health and Nutrition Examination Survey (NHANES) 2003–2006. *Crit Rev Food Sci Nutr* 2010;50(3):228–58. doi: 10.1080/10408391003626223.
8. Bowman SA. Diets of individuals based on energy intakes from added sugars. *Family Economics and Nutrition Review* 1999;12(2):31–8.
9. Murphy S, Johnson R. The scientific basis of recent US guidance on sugars intake. *Am. J. Clin. Nutr.* 2003;78:827S-833S.
10. Welsh JA, Sharma AJ, Grellinger L, et al. Consumption of added sugars is decreasing in the United States. *Am J Clin Nutr* 2011;94:726–34. doi: 10.3945/ajcn.111.018366.
11. Yang Q, Zhang Z, Gregg EW, et al. Added Sugar Intake and Cardiovascular Diseases Mortality Among US Adults. *JAMA Intern Med.* 2014 Apr;174(4):516-24. doi:10.1001/jamainternmed.2013.13563.
12. United States Department of Agriculture. Economic Research Service. Sugar and Sweeteners Yearbook Tables. <<http://www.ers.usda.gov/data-products/sugar-and-sweeteners-yearbook-tables.aspx>> (accessed August 2015).
13. National Health and Nutrition Examination Survey <http://wwwn.cdc.gov/nchs/nhanes/search/nhanes09_10.aspx> (accessed August 2015).
14. Johnson CL, Paulose-Ram R, Ogden CL, et al. National Health and Nutrition Examination Survey: Analytic guidelines, 1999–2010. National Center for Health Statistics. *Vital Health Stat* 2 (161). 2013.

- http://www.cdc.gov/nchs/data/series/sr_02/sr02_161.pdf (accessed August 2015)
15. NHANES. MEC In-Person Dietary Interviewers Procedures Manual. January 2009.
<http://www.cdc.gov/nchs/data/nhanes/nhanes_09_10/DietaryInterviewers_Inperson.pdf> (accessed August 2015)
16. NHANES. Phone Follow-Up Dietary Interviewer Procedures Manual. September 2009.
<http://www.cdc.gov/nchs/data/nhanes/nhanes_09_10/Dietary_PFU_09.pdf> (accessed August 2015)
17. Moshfegh AJ, Rhodes DG, Baer DJ, et al. The USDA Automated Multiple-Pass Method reduces bias in the collection of energy intakes. *Am J Clin Nutr* 2008;88:324-332.
18. Blanton CA, Moshfegh AJ, Baer DJ, et al. The USDA Automated Multiple-Pass Method accurately estimates group total energy and nutrient intake. *J Nutr* 2006 Oct;136(10):2594-9.
19. Rumpler WV, Kramer M, Rhodes DG, et al. Identifying sources of reporting error using measured food intake. *Eur J Clin Nutr* 2008;62:544-52.
20. Automated Multiple-Pass Method. United States Department of Agriculture. Agriculture Research Service.
<http://www.ars.usda.gov/ba/bhnrc/fsrg> (accessed August 2015).
21. National Health and Nutrition Examination Survey. NHANES Response Rates and Population Totals. Response Rates.
<http://www.cdc.gov/nchs/nhanes/response_rates_CPS.htm> (accessed August 2015).
22. National Health and Nutrition Examination Survey. 2009 - 2010 Data Documentation, Codebook, and Frequencies.
http://www.cdc.gov/nchs/nhanes/nhanes2009-2010/DR1IFF_F.htm (accessed August 2015).
23. Monteiro CA, Levy RB, Claro RM, et al. A new classification of foods based on the extent and purpose of their processing. *Cad. Saúde Pública*, Nov 2010, vol.26, no.11, p.2039-2049.
24. Moubarac JC, Parra DC, Cannon G, et al. Food Classification Systems Based on Food Processing: significance and implications for policies and actions: a systematic literature review and assessment. *Curr Obes Rep* 2014;3:256-272.
25. Monteiro CA, Cannon G, Levy RB, et al. Ultra-processing and a new classification of foods. In: Neff R (ed) *Introduction to U.S. Food System. Public Health, Environment, and Equity*. San Francisco: Jossey Bass A Wiley Brand; 2015.
26. Monteiro CA. Nutrition and health. The issue is not food, nor nutrients, so much as processing. *Public Health Nutr*. 2009;12(5):729-731.

27. Ludwig DS. Technology, diet, and the burden of chronic disease. *JAMA* 2011;305:1352-1353.

28. Monteiro CA, Cannon G, Levy RB, et al. The Food System. Processing. The big issue for disease, good health, well-being. *World Nutrition* 2012; 3:527–569. www.wphna.org (accessed August 2015).

29. Moodie R, Stuckler D, Monteiro C, et al. Profits and pandemics: prevention of harmful effects of tobacco, alcohol, and ultra-processed food and drink industries. *Lancet*. 2013 Feb 23;381(9867):670-9. doi: 10.1016/S0140-6736(12)62089-3.

30. Ahuja JKA, Montville JB, Omolewa-Tomobi G, Heendeniya KY, Martin CL, Steinfeldt LC, Anand J, Adler ME, LaComb RP, and Moshfegh AJ. 2012. USDA Food and Nutrient Database for Dietary Studies, 5.0. U.S. Department of Agriculture, Agricultural Research Service, Food Surveys Research Group, Beltsville, MD.

31. U.S. Department of Agriculture, Agricultural Research Service. 2011. USDA National Nutrient Database for Standard Reference, Release 24. Nutrient Data Laboratory Home Page. <<http://www.ars.usda.gov/ba/bhnrc/ndl>> (accessed August 2015).

32. Bowman SA, Clemens JC, Thoeig RC, Friday JE, Shimizu M, and Moshfegh AJ. 2013. Food Patterns Equivalents Database 2009-10: Methodology and User Guide [Online]. Food Surveys Research Group, Beltsville Human Nutrition Research Center, Agricultural Research Service, U.S. Department of Agriculture, Beltsville, Maryland. <<http://www.ars.usda.gov/ba/bhnrc/fsrg>> (accessed June 2013).

33. Louzada MLC, Martins APB, Canella DC, et al. Ultra-processed foods and dietary quality in Brazil. *Rev. Saúde Pública São Paulo* 2015;vol.49; Epub July 10, 2015. doi:10.1590/S0034-8910.2015049006132


34. Dietary Guidelines for the Brazilian Population 2014. 2nd Edition. Ministry of Health of Brazil.

35. Eicher-Miller HA, Fulgoni III VL, Keast DR. Contributions of Processed Foods to Dietary Intake in the US from 2003–2008: A Report of the Food and Nutrition Science Solutions Joint Task Force of the Academy of Nutrition and Dietetics, American Society for Nutrition, Institute of Food Technologists, and International Food Information Council. *J. Nutr.* 2012;142:2065S–2072S. doi: 10.3945/jn.112.164442.

36. International Food Information Council Foundation. Understanding our food communications tool kit, 2010. <<http://www.foodinsight.org/For-Professionals/Understanding-Our-Food/tabid/1398/Default.aspx>> (accessed August 2015)

37. Poti JM, Mendez MA, Ng SW, et al. Is the degree of food processing and convenience linked with the nutritional quality of foods purchased by US households? *Am J Clin Nutr.* 2015 Jun;101(6):1251-62. doi:10.3945/ajcn.114.100925.

38. Moubarac J-C, Martins APB, Claro RM, et al. Consumption of ultra-processed foods and likely impact on human health. Evidence from Canada. *Public Health Nutr.* 2013 Dec;16(12):2240-8. doi: 10.1017/S1368980012005009.
39. Monteiro CA, Levy RB, Claro RM, et al. Increasing consumption of ultra-processed foods and likely impact on human health: evidence from Brazil. *Public Health Nutr.* 2011 Jan;14(1):5-13. doi:10.1017/S1368980010003241.
40. Crovetto M M, Uauy R, Martins AP, et al. Household availability of ready-to-consume food and drink products in Chile: impact on nutritional quality of the diet. *Rev Med Chil.* 2014 Jul; 142(7):850-8. doi:10.4067/S0034-98872014000700005.
41. Monteiro CA, Moubarac JC, Cannon G, et al. Ultra-processed products are becoming dominant in the global food system. *Obes Rev.* 2013 Nov;14 Suppl2:21-8. doi:10.1111/obr.12107.
42. Canella DS, Levy RB, Martins APB, et al. Ultra-processed food products and obesity in Brazilian households (2008-2009). *PLoS One.* 2014 Mar 25;9(3):e92752. doi:10.1371/journal.pone.0092752. eCollection 2014.
43. Bingham S, Luben R, Welch A, et al. Epidemiologic assessment of sugars consumption using biomarkers: comparisons of obese and nonobese individuals in the European Prospective Investigation of Cancer Norfolk. *Cancer Epidemiol Biomarkers Prev* 2007; 16(8):1651-4.
44. Lafay L, Mennen L, Basdevant A, et al. Does energy intake underreporting involve all kinds of food or only specific food items? Results from the Fleurbaix Laventie Ville Sante (FLVS) study. *Int J Obes Relat Metab Disord.* 2000; 24:1500-6.
45. Pryer JA, Vrijheid M, Nichols R, et al. Who are the "low energy reporters" in the dietary and nutritional survey of British adults? *Int J Epidemiol.* 1997; 26: 146-54.



59x43mm (300 x 300 DPI)

Online Supplementary Material

Food classification according to processing

Food items were initially classified into four groups shown in Table 1. This was accomplished by taking into account, the following three variables from the NHANES recall databases: "Main Food Description", "Additional Food Description" and "SR Code Description". Thereafter, the food item classification was modified, if necessary, taking two variables into account: "Combination Food Type" and "Source of food". Thus, most "Frozen meals" or "Lunchables" or food items consumed in "Restaurant fast food/pizza" or acquired at a "Vending machine", were classified as ultra-processed foods.

As explained in the Subjects and Methods section, when Food Codes were judged to be a handmade recipe, the classification was applied to the underlying ingredients (SR Codes), to enable a more precise food item classification (1).

It must be noted, however, that SR Codes and their proportions are not necessarily the ingredients and proportions consumed by the participant. One of the reasons is that links between FNDDS 5.0 and SR24 were developed to estimate the nutrient content of a Food Code and not the ingredient intake (2). Furthermore, when assigning SR Codes to a Food Code the individual-specific variable "Modification Code" ("adjustments to predefined recipe ingredients that reflect more closely the food as described by the respondent" (2)) was not taken into account, as manual changes would have had been necessary to do so.

Absence of data or discrepancies regarding degree of processing were solved opting for the lesser degree of processing (conservative criterion), which could have led to a slight underestimation of ultra-processed food consumption.

Online Supplementary Material

We classified homemade recipes with unknown ingredients based on expected principal ingredients, which could slightly underestimate ultra-processed food consumption.

Regarding bread, the classification distinguishes between handmade bread (either homemade or made in restaurants or artisanal bakeries), and industrial bread (made in industrial bakeries or factories), either processed (when made only of ingredients used in the making of handmade breads -flour, yeast, water, salt, and, sometimes, walnuts, dried fruits and other whole foods-) or ultra-processed (when adding substances not commonly used in the making of handmade breads -such as hydrogenated fat, sugars, starches, and additives). In our study, because of the large amount of industrial breads with unknown ingredients (approximately 3.7% of all industrial bread had fully known ingredients) and the very low consumption of processed breads when ingredients were reported (approximately 2.3% of industrial breads were processed), we ended up classifying all industrial bread as ultra-processed foods. This could slightly overestimate ultra-processed food consumption.

Assessing energy and added sugar contents

For some handmade recipes, the sum of the “calorie intake per SR Code” (calculated by us) of all underlying SR Codes did not add up exactly to the “calorie intake per Food Code” (provided by NHANES). In these cases, the “final calorie intake per SR code” was calculated as follows:

$$\text{Final calorie intake per SR code} = \text{NHANES Calorie intake per Food Code} * \left(\frac{\text{Calculated Calorie intake per SR code}}{\sum_{n=1}^{\infty} \text{Calculated Calorie intake per SR Code}} \right)$$

The same was done for added sugars:

Online Supplementary Material

$$\text{Final added sugars intake per SR code} = \text{Added sugars intake per Food Code} * \left(\frac{\text{Added sugars intake per SR code}}{\sum_{n=1}^{\infty} \text{Added sugars intake per SR code}} \right)$$

where n = each of the Food Code underlying SR Codes

For peer review only

Online Supplementary Material

Supplementary Table 1. NOVA food classification based on the extent and purpose of industrial processing (adapted from 3,4)

Food groups and definition	Examples
<p>1 Unprocessed or minimally processed foods</p> <p>Unprocessed foods are those obtained directly from plants or animals (such as green leaves and fruits, or eggs and milk) and purchased for consumption without having undergone any alteration following their removal from nature. Minimally processed foods are natural foods that have been submitted to cleaning, removal of inedible or unwanted parts, fractioning, grinding, drying, fermentation, pasteurisation, cooling, freezing, or other processes which do not add substances to the original food. Purpose of minimum processes is to preserve foods and make it possible to store them and, sometimes, also to decrease stages of food preparation (cleaning and removing inedible parts) or facilitate their digestion, or render them more palatable (grinding or fermentation).</p>	<p>Natural, packaged, cut, chilled or frozen vegetables, fruits, potatoes, cassava, and other roots and tubers; bulk or packaged white, parboiled and wholegrain rice; whole or separated corn; grains of wheat and other cereals that are dried, polished, or ground as grits or flour; dried or fresh pasta made from wheat flour and water; all types of beans; lentils, chickpeas, and other legumes; dried fruits, fruit juices fresh or pasteurized without added sugar or other substances; nuts, peanuts, and other oilseeds without salt or sugar; fresh and dried mushrooms and other fungi; fresh and dried herbs and spices; fresh, frozen, dried beef, pork, poultry and other meat and fish; pasteurized, 'long-life' and powdered milk; fresh and dried eggs, yoghurt without sugar; and tea, herbal infusions, coffee, and tap, spring and mineral water.</p>
<p>2 Processed culinary ingredients</p> <p>These are substances extracted from natural foods or from nature itself by processes such as pressing, grinding, crushing, pulverising, and refining. Purpose of processing here is to obtain ingredients used in homes and restaurants kitchens to season and cook natural or minimally processed foods and to create with them varied and enjoyable dishes such as soups and broths, salads, rice and beans dishes, grilled or roasted vegetables and meat, and homemade breads, pies, cakes, and desserts.</p>	<p>Plant oils; coconut and animal fats (including butter and lard); table sugar, maple syrup (100%), molasses and honey; and table salt.</p>
<p>3 Processed foods</p> <p>These are relatively simple products manufactured essentially with the addition of salt or sugar or other substance of common culinary use, such as oil or vinegar, to natural or minimally processed foods. Purpose here is to prolong duration of foods and modify their palatability.</p> <p>If alcoholic beverages should be classified, drinks produced by the fermentation of group 1 food items such as wine, beer and cider will be classified in this group.</p>	<p>Canned and bottled vegetables, legumes or fruits; salted nuts or seeds; salted, smoked or cured meat or fish; canned sardine and tuna; cheeses, and breads made of wheat flour, yeast, water, and salt.</p>
<p>4 Ultra-processed foods</p> <p>These are food and drink products whose manufacture involves several stages and various processing techniques and ingredients, many of which are used exclusively by industry. Purpose of processing here is to create durable, accessible, convenient, and highly palatable, ready-to-drink, ready-to-eat, or ready-to-heat products typically consumed as snacks or desserts or as fast meals which replace dishes prepared from scratch.</p>	<p>Confectionery, soft drinks, sweetened juices and dairy drinks, powders for juices, sausages, chicken and fish nuggets or sticks and other pre-prepared frozen dishes, dried products such as cake mix, powdered soup, instant noodles, ready-seasonings, and an infinity of new products including packaged snacks, morning cereals, cereal bars, and 'energy' drinks. Sugar substitutes, sweeteners and all syrups (excluding 100% maple syrup). Breads and baked goods become ultra-processed products when, in addition to wheat flour,</p>

Online Supplementary Material

If alcoholic beverages should be classified, drinks produced by fermentation of group 1 food items followed by distillation and eventual addition of sugars or other substances, such as rum, whiskey, vodka, gin, and liqueurs, will be classified in this group.	yeast, water, and salt, their ingredients include substances such as hydrogenated vegetable fat, sugar, starch, whey, emulsifiers, and other additives.
--	---

For peer review only

Online Supplementary Material

Supplementary Table 2. Percentage of participants with more than 10% of total energy intake from added sugars, by demographic subgroups, according to quintiles of the dietary contribution of ultra-processed foods. US population aged 1 + years (NHANES 2009-2010)

		Quintiles of the dietary contribution of ultra-processed foods (% of total energy intake)				
		1st (n=1,937)	2nd (n=1,888)	3rd (n=1,814)	4th (n=1,779)	5th (n=1,899)
Gender	Men (n=4,634)	24.5	48.6	61.7	78.1	78.5*
	Women (n=4,683)	28.3	51.4	63.7	75.3	85.6*
Age (years)	1 to 5 (n=1,136)	17.0	45.5	61.3	71.0	84.3*
	6 to 11 (n=1,154)	33.1	54.0	76.5	82.4	90.0*
	12 to 19 (n=1,265)	39.9	62.8	66.2	83.0	87.1*
	20 to 39 (n=1,928)	28.8	53.4	64.1	79.7	82.7*
	40 to 59 (n=1,935)	26.0	49.1	59.6	71.7	76.7*
	60 and over (n=1,899)	22.8	43.6	58.6	71.9	71.1*
Race/ethnicity	Mexican American (n=2,064)	28.5	52.8	64.5	79.4	84.7*
	Other Hispanic (n=988)	41.7	59.4	62.2	80.1	85.2*
	Non-Hispanic White (n=3,984)	22.9	47.3	60.1	75.3	80.4*
	Non-Hispanic Black (n=1,726)	33.0	60.3	76.5	82.1	89.0*
	Other Race (including Multi-Racial) (n=555)	25.8	45.0	64.5	73.0	79.2*

Online Supplementary Material

Income to poverty*	0.00–1.30 (n=3,322)	31.1	58.8	72.3	81.0	86.5*
	>1.30–3.50 (n=3,062)	26.4	50.0	67.1	77.4	84.9*
	>3.50 and above (n=2,100)	23.0	46.1	52.0	72.8	75.2*
Educational attainment	<12 years (n=2,669)	32.9	50.6	68.7	76.8	86.4*
	12 years (n=2,136)	29.3	56.2	66.0	81.8	83.7*
	>12 years (n=4,398)	23.4	47.7	59.1	74.2	79.9*

*Significant linear trend across quintiles ($P \leq 0.001$), both in unadjusted and Poisson models adjusted for sex, age group (1–5, 6–11, 12–19, 20–39, 40–59, 60 + years), race/ethnicity (Mexican-American, Other Hispanic, Non-Hispanic White, Non-Hispanic Black and Other Race - Including Multi-Racial-), ratio of family income to poverty (SNAP 0.00–1.30, >1.30–3.50, and >3.50 and over) and educational attainment (<12, 12 years and >12 years).

Online Supplementary Material

References (for Online Supplementary Material).

1. Fitt E, Mak TN, Stephen AM, Prynne C, Roberts C, Swan G and Farron-Wilson M. Disaggregating composite food codes in the UK National Diet and Nutrition Survey food composition databank. *European Journal of Clinical Nutrition* (2010) 64, S32–S36.

2. Ahuja JKA, Montville JB, Omolewa-Tomobi G, Heendeniya KY, Martin CL, Steinfeldt LC, Anand J, Adler ME, LaComb RP, and Moshfegh AJ. 2012. USDA Food and Nutrient Database for Dietary Studies, 5.0. U.S. Department of Agriculture, Agricultural Research Service, Food Surveys Research Group, Beltsville, MD.

3. Moubarac JC, Parra DC, Cannon G, Monteiro CA. Food Classification Systems Based on Food Processing: significance and implications for policies and actions: a systematic literature review and assessment. *Curr Obes Rep* 2014; 3: 256-272.

4. Monteiro CA, Cannon G, Levy RB, Claro RM, Moubarac J-C. Ultra-processing and a new classification of foods. In: Neff R (ed) *Introduction to U.S. Food System. Public Health, Environment, and Equity*. San Francisco: Jossey Bass A Wiley Brand; 2015.

STROBE Statement—checklist of items that should be included in reports of observational studies

	Item No	Recommendation
Title and abstract	1	(a) Indicate the study's design with a commonly used term in the title or the abstract (Cross-sectional study; p.2) (b) Provide in the abstract an informative and balanced summary of what was done and what was found (p.2-3)
Introduction		
Background/rationale	2	Explain the scientific background and rationale for the investigation being reported (p.4-5)
Objectives	3	State specific objectives, including any prespecified hypotheses (p.5)
Methods		
Study design	4	Present key elements of study design early in the paper (p.6)
Setting	5	Describe the setting, locations, and relevant dates, including periods of recruitment, exposure, follow-up, and data collection (p.6-7)
Participants	6	(a) Cohort study—Give the eligibility criteria, and the sources and methods of selection of participants. Describe methods of follow-up (nap) Case-control study—Give the eligibility criteria, and the sources and methods of case ascertainment and control selection. Give the rationale for the choice of cases and controls (nap) Cross-sectional study—Give the eligibility criteria, and the sources and methods of selection of participants (p.6-7) (b) Cohort study—For matched studies, give matching criteria and number of exposed and unexposed (nap) Case-control study—For matched studies, give matching criteria and the number of controls per case (nap)
Variables	7	Clearly define all outcomes, exposures, predictors, potential confounders, and effect modifiers. Give diagnostic criteria, if applicable (p.7-10)
Data sources/measurement	8*	For each variable of interest, give sources of data and details of methods of assessment (measurement). Describe comparability of assessment methods if there is more than one group (p.7-10)
Bias	9	Describe any efforts to address potential sources of bias (lines 60-61)
Study size	10	Explain how the study size was arrived at (lines 54-60; lines 134-136)
Quantitative variables	11	Explain how quantitative variables were handled in the analyses. If applicable, describe which groupings were chosen and why (p.8-10)
Statistical methods	12	(a) Describe all statistical methods, including those used to control for confounding (p.9-10) (b) Describe any methods used to examine subgroups and interactions (p.9-10) (c) Explain how missing data were addressed (lines 134-136) (d) Cohort study—If applicable, explain how loss to follow-up was addressed (nap) Case-control study—If applicable, explain how matching of cases and controls was addressed (nap) Cross-sectional study—If applicable, describe analytical methods taking account of sampling strategy (lines 137-142) (e) Describe any sensitivity analyses (nap)

Continued on next page

Results		
Participants	13*	(a) Report numbers of individuals at each stage of study—eg numbers potentially eligible, examined for eligibility, confirmed eligible, included in the study, completing follow-up, and analysed (lines 54-61, lines 134-136) (b) Give reasons for non-participation at each stage (lines 54-61, lines 134-136) (c) Consider use of a flow diagram (nap)
Descriptive data	14*	(a) Give characteristics of study participants (eg demographic, clinical, social) and information on exposures and potential confounders (lines 147-158; Table 1) (b) Indicate number of participants with missing data for each variable of interest (lines 134-136) (c) Cohort study—Summarise follow-up time (eg, average and total amount) (nap)
Outcome data	15*	Cohort study—Report numbers of outcome events or summary measures over time (nap) Case-control study—Report numbers in each exposure category, or summary measures of exposure (nap) Cross-sectional study—Report numbers of outcome events or summary measures (lines 180-193; Table 1)
Main results	16	(a) Give unadjusted estimates and, if applicable, confounder-adjusted estimates and their precision (eg, 95% confidence interval). Make clear which confounders were adjusted for and why they were included (lines 194-224; Figure 1; Table 2) (b) Report category boundaries when continuous variables were categorized (Table 2) (c) If relevant, consider translating estimates of relative risk into absolute risk for a meaningful time period (nap)
Other analyses	17	Report other analyses done—eg analyses of subgroups and interactions, and sensitivity analyses (lines 220-221; Supplementary Table 2)
Discussion		
Key results	18	Summarise key results with reference to study objectives (lines 233-247)
Limitations	19	Discuss limitations of the study, taking into account sources of potential bias or imprecision. Discuss both direction and magnitude of any potential bias (lines 296-308)
Interpretation	20	Give a cautious overall interpretation of results considering objectives, limitations, multiplicity of analyses, results from similar studies, and other relevant evidence (lines 309-314)
Generalisability	21	Discuss the generalisability (external validity) of the study results (lines 265-288)
Other information		
Funding	22	Give the source of funding and the role of the funders for the present study and, if applicable, for the original study on which the present article is based (nap)

*Give information separately for cases and controls in case-control studies and, if applicable, for exposed and unexposed groups in cohort and cross-sectional studies.

Note: An Explanation and Elaboration article discusses each checklist item and gives methodological background and published examples of transparent reporting. The STROBE checklist is best used in conjunction with this article (freely available on the Web sites of PLoS Medicine at <http://www.plosmedicine.org/>, Annals of Internal Medicine at <http://www.annals.org/>, and Epidemiology at <http://www.epidem.com/>). Information on the STROBE Initiative is available at www.strobe-statement.org.